

VESTAL INSURE

Germicidal Detergent

EPA Reg. No. 1043-78

Germicide • Deodorizer • Fungicide • Virucide* • Cleaner

Active Ingredients:

Octyl decyl dimethyl ammonium chloride	2.90
Dioctyl dimethyl ammonium chloride	1.45
Didecyl dimethyl ammonium chloride	1.45
Alkyl (50% C ₁₂ , 40% C ₁₄ , 10% C ₁₆) dimethyl benzyl ammonium chloride	1.9

Inert Ingredients: 92.90

KEEP OUT REACH OF CHILDREN

DANGER:

Statement of practical treatment

In case of eye contact, flush with water for 15 minutes. Get medical attention. In case of skin contact, flush with water. See side panel for additional precautionary statements.

Contents: One U.S. Gallon

VESTAL INSURE

A One-Step Germicidal Detergent for Health Care Institutions and Industrial Use.

Vestal Insure is a specially formulated concentrated hardwater effective, phosphate free germicidal detergent. To use, simply add one ounce of Vestal Insure to each pre-measured gallon of water, gently stir to make a uniform solution and it is ready to go to work.

The fine detergents in Vestal Insure aid in cleaning while it simultaneously deodorizes and disinfects. It may be used for cleaning and disinfection of all washable hard non-porous surfaces such as floors, walls, woodwork, bathroom fixtures, equipment, and furniture. It is intended for use only in institutions such as hospitals, nursing homes, schools, industrial plants, medical and dental offices, veterinary clinics, and other public areas.

Use it for cleaning and disinfecting glass, chrome, stainless steel, porcelain, tile, washable painted or varnished surfaces, as well as resilient vinyl, linoleum, asphalt, rubber or terrazzo floors.

Germicidal: Passes A O A C Germicidal Use Dilution Method (S. aureus, S. choleraesuis, P. aeruginosa) in 400 ppm hard water, 1:128 (1 oz./1 gallon) in the presence of 5% organic soil (serum), 10 minutes at 20°C.

Fungicidal: Passes A O A C Fungicidal Test (Trichoglyphes) when diluted with 400 ppm hard water, 1:128 in the presence of 5% added organic soil (serum), 10 minutes at 20°C.

Virucidal: Passes Virucidal-Quantitation Test (Influenza A, Herpes Simplex, Vaccinia) when diluted with 400 ppm hard water, 1:128 (1 oz./1 gallon) in the presence of organic soil (serum), 10 minutes at 20°C.



Vestal Laboratories
A Division of Chemed Corp.
St. Louis, MO 63110

EPA Est. No. 1043-MO-1
6402 97A-100

36



Precautionary Statements

Hazards to Humans and Domestic Animals:

DANGER

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide Disposal: This sanitizer, its solutions or rinsings from empty containers should be disposed of in a toilet or service sink served by a sanitary sewer, in a land fill approved for pesticides or buried in a safe place away from water supplies.

Container Disposal: Triple rinse and deposit in a waste container for incineration or burial in a land fill approved for pesticide containers, or bury in a safe place.

General: Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning if frozen, thaw and remix before use.



VESTAL LABORATORIES, A Division of Chemed Corporation • St. Louis, MO 63110

ACCEPTED
MAY 12 1981
Used under Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1043-80

Vestal PROCES

No-Rinse Sanitizer for Food Processing, Hospital, Industrial, School, Dairy and Other Facilities

EPA Reg. No. 1043-80

Active Ingredient
Alkyl (C₁₂, 50%; C₁₄, 40%; C₁₆, 10%) Dimethyl Benzyl Ammonium Chloride
Inert Ingredients

KEEP OUT OF REACH OF CHILDREN
DANGER

Statement of Practical Treatment

- In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. Call a physician. Remove and wash contaminated clothing before reuse.
- If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or water. Do not induce vomiting. Drink large quantities of water. Avoid alcohol. Call a physician immediately.
- **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of emetics. Watch for signs of respiratory depression, and convulsion may be noted.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Contents: 1 U.S. Gallon