Precautionary Statements

HAZARD TO HUMANS AND DOMESTIC ANIMALS [illegible] corrosive. Causes eye and skin damage. Do not get on skin, on clothing, or on clothing. Wear goggles or face shield and rubber gloves when handling. *Harmful if swallowed. Concentrate may be fatal if swallowed through the skin.*

STATEMENT OF PRACTICAL TREATMENT: In case of eye contact, immediately flush eyes with water for 15 minutes and get prompt medical attention. For skin contact, flush thoroughly with water. Remove and wash contaminated clothing before reuse. If swallowed, drink large quantity of egg whites, gelatin solution, or water. Avoid alcohol. Do not induce vomiting. Call physician immediately. NOTE TO PHYSICIAN: Prolonged mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CLEANING AND DISINFECTING: Add one-half ounce (15cc) to each measured gallon of water used. Always add this product to premeasured water. Gently mix for a uniform solution. Apply solution with a cloth, sponge, mop, or brush using normal cleaning methods. Thoroughly wet all surfaces to be cleaned, then remove excess solution with a wring-out applicator. Treated surfaces should remain wet for ten minutes. Remove gross filth mechanically by sweeping before cleaning begins. Dispose solution when it becomes dirty and replace with a fresh solution. This is a corrosive product. Do not mix other chemicals. Use only as directed.

ANIMAL HUSBANDRY, ANIMAL RESEARCH, POULTRY FACILITY CLEANING AND DISINFECTING: Remove or cover all food, medicine, beverages, manure and livestock. Remove all litter, manure and other debris from floors and walls; scrape if necessary. Empty all troughs, mixers, feeders and waterers. Saturate all surfaces with a 1:256 solution (1/2 ounce per gallon prepared as directed above). Scrub to loosen adhering soils. Allow to soak for 10 minutes, then wash all surfaces with a wring-out applicator. Flush troughs and waterers with potable water after disinfecting. Drain all surfaces, remove footbaths and allow to air-dry before use or rewarding animals.

CLEANING AND DISINFECTING FOOD PREPARATION AND PROCESSING FACILITIES AND EQUIPMENT: Cover or remove all food and packaging materials. Remove all grease soils. Saturate all surfaces with this use-solution (1/2 ounce per gallon prepared as directed above). Scrub to remove all soils. Allow to soak for 10 minutes. Thoroughly rinse all treated and adjacent surfaces with potable water.

†KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected *likelihood of soiling of inanimate surfaces/objects* with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

†SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (HUMAN IMMUNODEFICIENCY VIRUS OR AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

PERSONAL PROTECTION: Wear appropriate barrier protection such as latex gloves, gowns, masks or eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of 1:256 use-solution (1/2 ounce per gallon). Prepare and apply solution as directed in paragraph above.

CONTACT TIME: While the HIV-1 virus is inactivated in 30 seconds, use a 10 minute contact time for disinfection of all organisms on this label.

INFECTIOUS MATERIALS DISPOSAL: Blood and other body fluids should be decontaminated and disposed of according to local regulations for infectious materials.

Kills HIV-1 (AIDS) virus on inanimate surfaces†

Coverage NPD™

ONE-STEP GERMICIDAL DETERGENT
E.P.A. Reg. No 1043-60

Germicide • Fungicide • Virucide*
Detergent • Deodorizer • Concentrate

Active Ingredients	
Octyl decyl dimethyl ammonium chloride	4.80%
Dioctyl dimethyl ammonium chloride	2.30%
Didecyl dimethyl ammonium chloride	2.30%
Alkyl (50% C_{14}, 40% C_{12}, 10% C_{16}) dimethyl benzyl ammonium chloride	6.14%
Inert ingredients	84.46%

FOR AGRICULTURAL, INDUSTRIAL AND INSTITUTIONAL USE

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT
In case of eye contact, flush with water for 15 minutes. Get prompt medical attention. For skin contact, flush with water. *See side panel for additional precautionary statements.*

Contents: 30 U.S. Gallons (113.55 Liters)

This product is a ready-to-use, pre-cleaned, hard surface, germicidal detergent deodorizer. Contains quaternary ammonium compounds designed to combine cleaning and disinfecting in one operation. Cleans and disinfects in the presence of 5% organic soil and remains active when diluted with 400 ppm hard water.

It is intended for use on washable, hard, non-porous, environmental surfaces such as floors, walls, partitions, tools, carts, and other equipment in food processing, preparation and service facilities, in animal husbandry, poultry and food processing facilities, veterinary clinics, animal research facilities, and in other indoor areas where infectious contamination control is a necessity.

Use this use-solution for the cleaning, deodorization of inert surfaces such as stainless or hard-surface vinyl and asbestos, (polymer, masonry, glass, tile, varnished, painted or varnished surfaces, surfaces, as well as resilient vinyl, asphalt, linoleum, rubber, laminate or other commercial-type floors.

Germicide: Passes A.O.A.C. Germicidal Use-Dilution Method (S. aureus, S. choleraesuis, P. aeruginosa) when diluted with 400 ppm A.O.A.C. hard water or make a 1:256 (1/2 ounce per 1 gallon) solution in the presence of 5% organic soil (albumin), 10 minutes at 20°C.

It has also been effective in the A.O.A.C. Use-Dilution Test when diluted with 400 ppm A.O.A.C. hard water to make a 1:256 solution (1/2 ounce per 1 gallon of water) in the presence of 5% organic soil (albumin), 10 minutes at 20°C to inactivate

Staphylococcus aureus, ATCC 6538	Pseudomonas aeruginosa, ATCC 15442
Candida albicans, ATCC 10231	Salmonella choleraesuis, ATCC 10708
Streptococcus faecalis, ATCC 10541	Salmonella typhosa, ATCC 6539
Escherichia coli, ATCC 11229	Shigella dysenteriae, ATCC 12924
Trichophyton mentagrophytes, ATCC 9533	Shigella sonnei, ATCC 9290
Staphylococcus aureus, ATCC 6538	Streptococcus pyogenes, ATCC 19615
Staphylococcus aureus, ATCC 29213	Streptococcus pneumoniae, ATCC 6303
Escherichia coli, ATCC 25922	Streptococcus agalactiae, ATCC 13813
Pseudomonas aeruginosa, ATCC 15442	Streptococcus faecalis, ATCC 19433
Salmonella choleraesuis, ATCC 10708	Streptococcus pyogenes, ATCC 19615
Shigella dysenteriae, ATCC 12924	Streptococcus pneumoniae, ATCC 6303
Shigella sonnei, ATCC 9290	Streptococcus agalactiae, ATCC 13813
Streptococcus pyogenes, ATCC 19615	Streptococcus faecalis, ATCC 19433
Streptococcus pneumoniae, ATCC 6303	Streptococcus pyogenes, ATCC 19615
Streptococcus agalactiae, ATCC 13813	Streptococcus pneumoniae, ATCC 6303
Streptococcus faecalis, ATCC 19433