



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 29, 2018

Mike Ebers
Steris Corporation
P.O. Box 147
St. Louis, MO 63166-0147

Subject: PRIA Label Amendment – Revision of Precautionary Statements
Product Name: LpH® Master Product
EPA Registration Number: 1043-91
Application Date: Sept. 11, 2017
Decision Number: 533042

Dear Mr. Ebers:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 1043-91
Decision No. 533042

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Srinivas by phone at (703) 308-6354, or via email at gowda.srinivas@epa.gov.

Sincerely,



Jacqueline Hardy
Product Manager 34
Regulatory Management Branch II
Antimicrobials Division 7510P

Enclosure: Stamped Accepted Label

ACCEPTED

01/29/2018

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 1043-91

LpH[®]

Master Product

STERILE

ONE-STEP CLEANER DISINFECTANT

Reorder Number:

Net Contents: ½ fl. oz., 1 fl. oz., 1 gal., 5 gal., 55 gal.

Germicide Fungicide Virucide* Tuberculocide Deodorizer Cleaner Detergent

When diluted with soft or hard water (400 ppm CaCO₃) even in the presence of organic soil (5% serum)

EPA REG. NO. 1043-91

EPA Est. No. 1043-MO-002
211-KS-001

FOR AGRICULTURAL, INDUSTRIAL AND INSTITUTIONAL USE

Active Ingredients:

o-phenylphenol..... 7.7%

p-tertiary amylphenol.....7.6%

Inert Ingredients:.....84.7%

Total 100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

FIRST AID	
If inhaled	<ul style="list-style-type: none"> •Move person to fresh air. •If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. •Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15-20 minutes. •Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> •Hold eye open and rinse slowly and gently with water for 15-20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. •Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> •Call poison control center or doctor immediately for treatment advice. •Have person sip a glass of water if able to swallow. •Do not induce vomiting unless told to do so by the poison control center or doctor. •Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN	
Probable mucosal damage may contraindicate the use of gastric lavage.	

STERILE UNLESS PACKAGING IS COMPROMISED

STERIS Corporation
7501 Page Avenue
St. Louis, Mo 63133 U.S.A.
800-548-4873
www.steris.com

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS: DANGER: Corrosive. Causes irreversible eye damage. Causes skin burns. Harmful if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Wear protective eyewear, protective clothing and gloves. Wash hands thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash clothing before reuse.

(5 gal. or greater package only) ENVIRONMENTAL HAZARD STATEMENT: Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

This state of the art, non-alkaline, phenolic germicidal detergent is a sterile, multi-purpose product specifically developed to be both hard water and serum effective in a concentrated system for use on hard, non-porous critical environment surfaces found in aseptic rooms and hoods; air locks; gowning; barrier isolation; process equipment and piping systems. The use-solution is designed to clean, disinfect and deodorize in a simple one step process. In addition the non-alkaline formula removes soap scum and hard water scale which can harbor bacteria. Use this product on washable hard non-porous environmental surfaces such as floors, walls, counters, tools, carts, and other equipment in food processing preparation and service facilities, in animal and avian husbandry, poultry and feed processing facilities, veterinary clinics, animal research facilities, pharmaceuticals, biotechnology, medical device and FDA regulated facilities.

GERMICIDAL-The 1:256 (one half ounce per gallon) solution passes the A.O.A.C. Germicidal Use-Dilution Test (*S. aureus*, *S. enterica*, *Ps. aeruginosa*) when diluted with 400 ppm A.O.A.C. hard water, in the presence of 5% organic soil (serum), 10 minutes at 20°C.

BROAD SPECTRUM DATA: The following organisms also pass the A.O.A.C. Use-Dilution Test in 400 ppm A.O.A.C. hard water at a dilution of 1:256 in the presence of 5% organic soil (serum), 10 minutes at 20°C.

Acinetobacter calcoaceticus, ATCC 19606
Aeromonas hydrophila, ATCC 7965
Bordetella avium, ATCC 35086
Campylobacter jejuni, ATCC 29428
Candida albicans, Clinical Isolate
Candida parapsilosis, Clinical Isolate
Citrobacter freundii, ATCC 8090
Enterobacter aerogenes, ATCC 13048
Enterobacter cloacae, ATCC 23355
Escherichia coli, ATCC 25922
Klebsiella pneumoniae, ATCC 13883
Listeria monocytogenes, ATCC 15313
Mycoplasma gallisepticum, ATCC 19610
Pasteurella multocida, ATCC 29977
Proteus mirabilis, Clinical Isolate
Proteus vulgaris, ATCC 13315

Pseudomonas aeruginosa, ATCC 27853
Salmonella enteritidis, ATCC 13076
Salmonella sp. serovar pullorum, ATCC 19945
Salmonella typhi, ATCC 6539
Salmonella typhimurium, ATCC 14028
Serratia marcescens, ATCC 8100
Shigella flexneri, ATCC 12022
Shigella sonnei, ATCC 25931
Staphylococcus aureus, ATCC 25923
Staphylococcus aureus, (MRSA),
MultiPLY (Methicillin)-Resistant
Clinical Isolate
Staphylococcus epidermidis, ATCC 12228
Streptococcus faecalis, ATCC 19433
Streptococcus pyogenes, ATCC 19615
Streptococcus suis, ATCC 43765

FUNGICIDAL: The 1:256 (one half ounce per gallon) solution passes the A.O.A.C. Fungicidal Test (T. Mentagrophytes) when diluted with 400 ppm A.O.A.C. hard water, in the presence of 5% organic soil (serum), 10 minutes at 20°C.

TUBERCULOCIDAL: The 1:256 (one half ounce per gallon) solution passes the A.O.A.C. Tuberculocidal Test (Mycobacterium tuberculosis, var. bovis (BCG)) when diluted with 400 ppm A.O.A.C. hard water, in the presence of 5% organic soil (serum), 10 minutes at 20°C.

***VIRUCIDAL:** A 1:256 (one half ounce per gallon) solution passes the Virucidal Assay (EPA Proposed Method) against the following on hard inanimate surfaces when diluted with 400 ppm A.O.A.C. hard water, in the presence of 5% organic soil (serum), in 10 minutes at 20°-25°C. Influenza A₂ [Japan] virus, Herpes Simplex Type 2 virus, Vaccinia virus, Adenovirus Type 2 virus, Avian Reovirus, Avian Influenza virus, Newcastle Disease virus, Canine Distemper virus, Pseudorabies virus, Infectious Bronchitis virus, Porcine Rotavirus, Infectious Laryngotracheitis virus, Transmissible Gastroenteritis virus, Marek's Disease virus.

A 1:64 (2 fl. oz. per gallon) solution passes the Virucidal Assay against Canine Parvovirus and Infectious Bursal Disease virus on hard inanimate surfaces when diluted with 400 ppm A.O.A.C. hard water, in the presence of 5% organic soil (serum) 10 minutes at 20° - 25°C.

†When tested by a Dilution Method, the HIV-1(AIDS) virus, with added 5% organic soil (serum), was completely inactivated by a 1:256 (one half ounce per gallon) solution in 400 ppm A.O.A.C. hard water in 60 seconds at 20°C. Although efficacy at 1 minute contact time has been shown to be adequate against HIV-1, this time would not be sufficient for other organisms. Use a 10-minute contact time for disinfection against all of the organisms claimed.

This product has demonstrated effectiveness against influenza A virus and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 influenza A virus.

or

Kills Pandemic 2009 H1N1 influenza A virus formerly called swine flu.

ODOR CONTROL: It eliminates many odors by killing odor-causing bacteria while simultaneously chemically neutralizing their odors.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CLEANING AND DISINFECTING WASHABLE, HARD, NON-PORUS SURFACES: To make solution, add one-half fl. oz (15 cc) to each pre-measured gallon of water and gently mix until solution is uniform. **NOTE:** Always add this product (concentrate) to pre-measured water. Apply solution with a cloth, sponge, coarse spray, mop or brush using normal cleaning methods. Allow treated surfaces to soak for 10 minutes. Then remove excess solution with a wrung-out applicator. As solution becomes dirty, discard and replace with a fresh one. Before cleaning begins, always sweep or dust mop area to be cleaned thoroughly, removing any gross filth. A properly prepared solution of this product intended for use as a hard surface disinfectant has a shelf life of 14 days (use life) when stored in a closed container such as a spray bottle. This is a complete product. Do not add other chemicals. If frozen, thaw and re-mix before use. Use only as directed.

ANIMAL HUSBANDRY, ANIMAL RESEARCH, POULTRY, POULTRY HOUSE FACILITY

CLEANING AND DISINFECTION: Do not use in milking stalls, milking parlors or milk houses. Remove or cover all food, feedstuffs, packaging materials and livestock. Remove all litter, manure and other debris from floors and walls; scrape if necessary. Empty all troughs, racks, feed and water appliances. Saturate all surfaces with a 1:256 solution (one half ounce per gallon prepared as directed above). Scrub to loosen adhering soils. Allow to soak for 10 minutes, then pick up excess with a wrung-out applicator. Flush feeders and waterers with potable water after disinfecting. Drain all surfaces, ventilate buildings and allow to air dry before use or rehousing animals.

FOGGING AS AN ADJUNCT TO REGULAR CLEANING AND DISINFECTING: Before fogging remove or cover any food or packaging material with waterproof coverings. This product may be used in fogging (wet misting) as an adjunct following regular cleaning and disinfecting procedures in research and manufacturing facilities, in poultry operations, hatcheries, swine operations, animal care facilities, animal research facilities, livestock operations, veterinary facilities, zoos and kennels. Use a 1:256 dilution (one-half ounce per gallon of water). Remove all animals and feed from areas to be fogged. Remove all litter and manure from surfaces or areas traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces. Fog the desired area at 32 to 64 ounces of use dilution per 1000 cubic feet using equipment with an automatic timer. Do not remain in treated areas; allow at least two hours after fogging is complete before reentering fogged area. Treated surfaces must be thoroughly scrubbed with LpH Master Product at a 1:256 dilution and rinsed with potable water before reuse. Do not rehouse livestock or employ equipment until treatment has dried.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (HUMAN IMMUNODEFICIENCY VIRUS OR AIDS VIRUS) OF SURFACES / OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Wear appropriate barrier protection such as latex gloves, gowns, masks and eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of a 1:256 use-solution (one-half ounce per 1 gallon). Prepare and apply solution as directed in paragraph above.

Contact Time: While the HIV-1 virus is inactivated in 1 minute, use a 10-minute contact time for disinfection of all organisms on this label.

Infectious Materials Disposal: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious disposal.

CLEANING AND DISINFECTING FOOD PREPARATION, PROCESSING FACILITIES AND EQUIPMENT: Cover or remove all food and packaging materials. Remove all gross soils. Saturate all surfaces with the use-solution (one-half ounce per gallon prepared as directed above). Scrub to loosen all soils. Allow to soak for 10 minutes, then pick up excess solution with a wrung-out applicator. Thoroughly rinse all wetted and cleaned food contact surfaces with potable water.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not store near heat or open flame. If frozen, thaw and remix before use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

(For Packet:) Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

(For ≤5 gal:) Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(For > 5 gal:) Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining

contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

GENERAL: Consult Federal, State, or Local disposal authorities for approved alternative procedures such as limited open burning.