

ACCEPTED

MAY 6 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 1057-28

DOLGE WARFARIN

MIXED BAIT READY TO USE

For controlling rats and house mice

WARFARIN BAIT will not give effective control when eaten by rats and house mice in a single feeding. It must be eaten at several feedings (usually 5 or more) on successive days, with no period longer than 48 hours between feedings.

DIRECTIONS: Bait is ready-mixed. DO NOT add anything. Use at least 4 ounces of bait per bait location. An uninterrupted supply should be maintained for at least 10 days. Place bait in dry locations frequented by rats and mice. Place bait in shallow containers or preferably in protected feeding stations. Put a cup full or more of bait in each. Inspect stations daily and add bait as needed, increasing the amount when containers are emptied over night. Continue as long as any bait is taken, which may be from two to four weeks.

For Roof Rats put bait at GROUND FLOOR AND TOP FLOOR OR ATTIC levels. For Norway rats put bait at or near ground level, and at burrows and harborages. For house mice place tablespoon amounts of bait at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

CAUTION

KEEP OUT OF REACH OF CHILDREN

This bait would be dangerous to humans, domestic animals, pets or wildlife if taken in a large single dose, or in repeated small doses over a long period of several days. Baits should be placed where small children, irresponsible persons and domestic animals, pets or wildlife cannot get at them. CAREFUL placement of bait is therefore required. Use protected feeding stations wherever a hazard exists.

FIRST AID: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately. (Note to physicians: When a human has been known to have ingested warfarin bait, blood transfusions combined with intravenous injection and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol.) Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

ACTIVE INGREDIENT: Warfarin 3-(alpha-acetonylbenzyl)-	
4-hydroxycoumarin	0.025%
INERT INGREDIENTS:	99.975%
Total	100.00%

Westport  Conn.

5 POUNDS NET WEIGHT

WAR472-A

E.P.A. REG. NO. 1057-28