

WARNING: KEEP OUT OF REACH OF CHILDREN - SEE ANTICOLL AND OTHER WARNINGS ON LABEL

Consult your State Agricultural Extension Service for additional information consistent with label directions and cautions.

WARNINGS

May be fatal if swallowed. Do not get in eyes, on skin, or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes get medical attention. Do not breathe vapor or dust. May cause poisoning by inhalation or skin absorption. Wash thoroughly with soap and water after handling or after soiling or smudging; wear clean clothing. During commercial application of exposure in mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U.S.D.A. for Heptachlor protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate feed or feed stuff. Use other than directed may contain toxic crops.

This product is highly flammable. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label.

Do not reuse containers. Destroy by burning, with venting or burning. Stay away from smoke or fumes.

Do not dispose of any unused material in such a manner that contact with humans or beneficial animals may occur.

ANTIBIOTIC: In case of spillage on skin, wash immediately with soap and water. Inhaled—Call a physician immediately. Unless otherwise indicated, induce vomiting by inducing a finger into the throat or by giving vomit salt water (one half teaspoon of salt to a glass of water orally).

NON-VOLATILE PESTICIDE

Follow directions carefully. Soil insecticides such as those listed here are not effective against all soil insects. The degree of control depends on the soil conditions, weather, and crop conditions, and other influencing factors in the use of the product. For the control of the *S. litig.* soil insecticide is not a soil insecticide, but a soil insecticide (including a variety of *mercurial*), with a degree of soil insecticide effect of use. Do not use in soil to which may require soil insecticide on this label.

ACTIVE INGREDIENTS (by weight)	
Heptachlor*	20.00%
Related compounds	7.76%
INERT INGREDIENTS (by weight)	72.24%
TOTAL	100.00%
*Heptachlor is 4.7 Methyl Chloride	

CONTROL OF SOIL INSECTS ON CROPS

Soil Insect Complex (Non-resistant Corn Rootworms, Seed Corn Beetle, Soil Corn Maggot, Wireworm, Green Coleaspis, Cutworms): For broadcast application, apply 7 1/2 to 10 pounds of 70% Heptachlor Granules per acre, preferably 10 to 14 days before planting. Disc or work into top 3 to 4 inches of soil immediately after application. For row treatment, apply 5 pounds of 70% Heptachlor Granules per acre. Broadcast treatment is preferred where Cutworms and/or Wireworm are special problems. See specific recommendations for Wireworm and Cutworm control.

COLLEOPTERIFORM (Non-resistant species and when such use is required for Coleoptera spp.): Use 2 to 5 pounds of 70% Heptachlor Granules per acre as a row treatment. For broadcast treatment use 5 to 7 1/2 pounds of 70% Heptachlor Granules per acre. Mix into top 3 to 4 inches of soil immediately after broadcast application. In Missouri and possibly other states—check state recommendations—use 5 pounds row, and 7 1/2 pounds broadcast.

CUTWORMS: Apply 5 to 10 pounds of 70% Heptachlor Granules per acre as a broadcast treatment, preferably 10 to 14 days before planting. Mix into top 3 to 4 inches immediately after application. In some states a minimum dosage rate of 7 1/2 pounds is recommended.

WIREWORMS: Use 7 1/2 to 15 pounds of 70% Heptachlor Granules per acre as a preplant broadcast soil application, preferably 10 to 14 days before planting. Thoroughly work into top 3 to 4 inches of soil. Use 15 pounds dosage on potato hills or for high population of large worms.

WHITE GRUBS: Use 10 to 15 pounds of 70% Heptachlor Granules per acre as a broadcast treatment, preferably 10 to 14 days before planting and immediately before.

SEED CORN BEETLES AND SOIL CORN MAGGOTS: Use 7 1/2 to 10 pounds of 70% Heptachlor Granules per acre as a preplant broadcast treatment, preferably applied 10 to 14 days before planting. Disc or work into top 3 to 4 inches of soil immediately after application. A broadcast or row treatment apply 2 1/2 to 5 pounds in the row at planting time.