

AC
J
UNDE
FUNG
FOR
D U

E.W.T. plus

CONTROLS BROAD-LEAVED WEEDS WILL NOT HARM USUAL LAWN GRASSES

DIRECTIONS

Time of application: Apply when weeds are growing vigorously. Evidence of discoloration may result if applied in very hot weather (over 90° F). Do not apply in drought.

Rate of Application: 4 pints per acre. Must be mixed with water before use. For best results use 1½ fl. oz. diluted in 2 gallons of water per 1000 sq. ft. on smaller areas; 4 pints diluted in 30-40 gallons of water per acre for larger areas.

Mixing: To insure uniform mixing fill half the spray tank with water; then add required amount of E.W.T. PLUS. Mix thoroughly, adding remainder of water as required.

Application: Always use equipment suitable for uniform application. A boom-type sprayer is recommended. For absorption of the chemical by weeds, avoid cutting or mowing turf for 2-3 days before or after spraying. Avoid watering for 24 hours after spraying.

Full effects of E.W.T. PLUS are not visible for about 2-3 weeks.

WEEDS CONTROLLED

Susceptible: clover (*Trifolium* spp.), common chickweed (*Stellaria media*), dandelion (*Taraxacum* spp.), lambsquarters (*Chenopodium album*), mouse-ear chickweed (*Cerastium vulgatum*), pigweed (*Amaranthus* spp.), plantain (*Plantago* spp.), ragweed (*Ambrosia* spp.).

Moderately susceptible*: black medic (*Medicago lupulina*), dock (*Rumex* spp.), English daisy (*Bellis perennis*), knotweed (*Polygonum* spp.), mallow (*Malva* spp.), purslane (*Portulaca oleracea*), wood sorrel (*Oxalis* spp.).

* For best results apply E.W.T. PLUS during early stages of growth.

E.W.T. PLUS is a stable solution containing a mixture of phenoxy acids equivalent to 48 oz. of acid per gallon. It is non-flammable and non-corrosive. It may be used on ornamental turf grasses such as bluegrasses, fescues, Bermuda and bent grasses on fairways.

ACTIVE INGREDIENTS: Diethanolamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid 31.00%

Diethanolamine salt of 2,4-dichlorophenoxyacetic acid 15.35%

INERT INGREDIENTS: 53.65%

100.00%


Equivalent to 20.9% of 2-(2-methyl-4-chlorophenoxy) propionic acid and 10.45% of 2,4-dichlorophenoxyacetic acid or 2 lbs. of 2-(2-methyl-4-chlorophenoxy) propionic acid and 1 lb. of 2,4-dichlorophenoxyacetic acid per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS



Harmful if swallowed. Avoid contact and other parts of body contaminated. This material is active at low concentrations. Do not spray flower beds, borders or drift may damage neighboring crops, beans, tomatoes, peas, cotton, grapes. Ponds, streams and rivers must not be washed. Wash equipment thoroughly with water out, fill with water and leave overnight. Product. Traces can harm susceptible crops. Store containers, tightly closed, away from other materials applied on crops. Protect from freezing. Used containers should be disposed of properly. E.W.T. PLUS is not recommended for use on these species. Bent grasses are more susceptible at fairway cutting height. Do not use on flowers and vegetables. Do not use on



E.W.T. plus

**CONTROLS BROAD-LEAVED WEEDS
WILL NOT HARM USUAL LAWN GRASSES**

E.W.T. PLUS is a stable solution containing a mixture of phenoxy acids equivalent to 48 oz. of acid per gallon. It is non-flammable and non-corrosive. It may be used on ornamental turf grasses such as bluegrasses, fescues, Bermuda and bent grasses on fairways.

ACTIVE INGREDIENTS: Diethanolamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	31.00%
Diethanolamine salt of 2,4-dichlorophenoxyacetic acid	15.35%
INERT INGREDIENTS:	53.65%
	100.00%

Equivalent to 20.9% of 2-(2-methyl-4-chlorophenoxy) propionic acid and 10.45% of 2,4-dichlorophenoxyacetic acid or 2 lbs. of 2-(2-methyl-4-chlorophenoxy) propionic acid and 1 lb. of 2,4-dichlorophenoxyacetic acid per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS

Westport



Conn.

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes and clothing. After using, wash hands and other parts of body contaminated with concentrate, spray or washings.

This material is active at low concentrations.

Do not spray flower beds, borders or shrubbery. Do not spray in windy weather as the spray drift may damage neighboring crops. Avoid spray drift to susceptible plants: it may injure beans, tomatoes, peas, cotton, grapes, ornamentals, etc. Coarse sprays are less likely to drift.

Ponds, streams and rivers must not be contaminated with the concentrate, spray or washings.

Wash equipment thoroughly with water and Cidet detergent immediately after use. Spray out, fill with water and leave overnight. Spray out again before storing or using for another product. Traces can harm susceptible crops sprayed later.

Store containers, tightly closed, away from food, water supplies, and seeds, fertilizers and other materials applied on crops.

Protect from freezing.

Used containers should be disposed of safely.

E.W.T. PLUS is not recommended for St. Augustine and centipede grasses, as it can damage these species. Bent grasses are more susceptible to this product and should be sprayed at fairway cutting height. Do not use on newly seeded lawns or turf for six months after planting.

Avoid drift onto other plants and do not use first cuttings after treatment as a mulch for flowers and vegetables. Do not use on golf course greens.

EPA Reg. No. 1057-57

ly. Evidence of discoloration
ply in drought.

er before use. For best results
smaller areas; 4 pints diluted

er; then add required amount
quired.

ation. A boom-type sprayer is
id cutting or mowing turf for
fter spraying.

llaria media), dandelion (Tar-
r chickweed (Cerastium vulga-
ragweed (Ambrosia spp.).

ck (Rumex spp.), English daisy
ve spp.), purslane (Portulaca

with.

ACCEPTED
 Jun 24, 1977
 UNDER THE FEDERAL INSECTICIDE ACT
 PESTICIDE REGISTRATION NO. 1457-57

E.W.T. plus

**CONTROLS BROAD-LEAVED WEEDS
 WILL NOT HARM USUAL LAWN GRASSES**

E.W.T. PLUS is a stable solution containing a mixture of phenoxy acids equivalent to 48 oz of acid per gallon. It is non-flammable and non-corrosive. It may be used on ornamental turf grasses such as bluegrasses, fescues, bermuds and bent grasses on lawns.

ACTIVE INGREDIENTS: Diethanolamine salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	31.00
Diethanolamine salt of 2-(4-chlorophenoxy)acetic acid	15.35
INERT INGREDIENTS	53.65
	100.00

Equivalent to 20.9% of 2-(2-methyl-4-chlorophenoxy)propionic acid and 10.45% of 2-(4-chlorophenoxy)acetic acid or 2 lbs. of 2-(2-methyl-4-chlorophenoxy)propionic acid and 1 lb. of 2-(4-chlorophenoxy)acetic acid per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN
 SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS

Westport



Conn.

CAUTION

Use only on lawns. Avoid contact with knicker and clothing. After using, wash hands and other parts of body contaminated with concentrate, spray or washings.

This material is a five percent concentration.

Do not spray flowers, beds, borders or shrubbery. Do not spray in windy weather as the spray drift may damage neighboring crops. Avoid spray drift to susceptible plants, it may injure beans, tomatoes, peas, cotton, grapes, or ornamentals, etc. Coarse sprays are less likely to drift. Birds, animals and man must not be contaminated with the concentrate, spray or washings.

Wash equipment thoroughly with water and clean detergent immediately after use. Spray container with water and leave overnight. Spray container before storing or using for another purpose. The product is harmful to susceptible crops sprayed later.

Store containers tightly closed away from food, water supplies, and seeds, fertilizers and other materials applied on crops.

Protect from freezing.

Used containers should be disposed of safely.

E.W.T. PLUS is not recommended for St. Augustine grass or bermuda grass, as it can damage these species. Bermuda grasses are more susceptible to this product and should be sprayed at maximum cutting height. Do not use on newly seeded lawns or turf for six months after planting. Avoid drift onto other plants and do not use first cuttings after treatment as a mulch for flowers and vegetables. Do not walk on treated grass.