

2-5 73  
1057-61

# MINTOL-128

FRAGRANT MINT DISINFECTANT

## GENERAL DIRECTIONS

For general industrial and institutional disinfection such as walls, floors, toilet seats, urinals, garbage receptacles, use one ounce of Mintol-128 per gallon of water. Always pour water on the disinfectant and stir thoroughly to a uniform milky emulsion.

All surfaces should be cleaned before using disinfectant. Thoroughly wet the surface with recommended dilution of the disinfectant. Allow solution to remain in contact with surface as long as possible, at least not less than 10 minutes. All food contact surfaces such as food carts must be rinsed with potable water for reuse.

## HOSPITALS

To disinfect and deodorize walls, floors and other surfaces, and for routine concurrent and terminal disinfection in all areas, mop or wipe with 1 ounce per gallon. Rinsing not required.

OPERATING ROOM — Mintol-128 is safe on any conductive flooring. It meets the requirements of the National Fire Protection Association. Use 1 ounce per gallon.

Phenol Coefficient A.O.A.C.: Staphylococcus aureus 13, Salmonella typhosa 12.

A.O.A.C. Use Dilutions: Staphylococcus aureus 1:128; Salmonella choleraesuis 1:128. Pseudomonas aeruginosa 1:128.

A.O.A.C. Fungicidal Test, Trichophyton interdigitale 1:256.

A.O.A.C. Tuberculocidal Test, Mycobacterium tuberculosis 1:128.

ACTIVE INGREDIENTS: Isopropanol, potassium ricinoleate, o-phenylphenol, o-benzyl p-chlorophenol, p-tertiary amylphenol, tetrasodium ethylenediaminetetraacetate, essential oils, propargyl alcohol ... 38.33%

INERT INGREDIENTS: Water, dyestuff ... 61.67%

TOTAL ... 100.00%

## DANGER

### KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles and face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink PROMPTLY a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Do not reuse empty drum. Return to drum conditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

Rinse empty 5- and 1-gallon containers thoroughly with water and discard them.

The C.B. DOLGE Co.

Westport



Conn.

MTL-273 A

E.P.A. Reg. No. 1057-61