

[1057-64]

0000000000  
0000000000

# DOLGE GENERAL PURPOSE AQUEOUS PYRETHRUM INSECTICIDE

## DIRECTIONS INDOOR APPLICATION

Use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacuum treated area and ventilate before reoccupying. For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

**SURFACE SPRAYING:** To control: *measles, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw toothed grain beetle, split or beetles, cigarette beetle, Drugstore beetles, meal worms, grain mites, and fleas.* Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray on the tops, sides, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery behind shelves and drawers. For silverfish spray bookcases. For ants spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control: *carpet beetles.* Spray edges of carpeting and under carpeting and rugs. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control: *bedbugs.* Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, and floors. Repeat treatment as needed.

To control: *fleas and brown dog ticks.* Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concentrated treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control: *larder moths.* Remove any infested material from storage. Thoroughly ventilate air for several hours, in contact if possible. Apply a rapid knockdown spray to all shelves, bins, and other storage, drawing the spray into a hole, crevice, and crevice. After the infested material is treated, lightly sweep or vacuum at least three feet from these areas. Repeat application at 30 day intervals if treated clothes are not stored in airtight containers.

A CONCENTRATED MICRO EMULSION

- CLEAR
- STABLE
- LEAVES NO OILY OR OBJECTIONABLE RESIDUE

EFFECTIVE, FAST ACTING INSECTICIDE.

FOR USE IN HOMES, INSTITUTIONS, RESTAURANTS,

FOOD PROCESSING PLANTS, DAIRIES, LIVESTOCK AND OUTDOORS FOR MOSQUITO CONTROL.

APPLY WITH CONVENTIONAL MECHANICAL OR COMPRESSED AIR EQUIPMENT.

## CAUTION

KEEP OUT OF REACH OF CHILDREN

See Additional Precautions On Right Hand Panel



**SPACE SPRAYING AND FOGGING:** Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Argentine grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths, direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space.

**OUTDOOR ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery of vegetation where mosquitoes may rest. Shrubbery and vegetation around treatment ponds, marsh, swam, ponds, and other areas may be treated. Application of this product to any body of water is prohibited. Apply at a rate of 1 1/2 gallons per acre (4 1/2 ounces per 1000 square feet).

Using a mounted compressed air spraying equipment such as Rotomist® models 510, 91 or 101 HT, traveling at 5 m.p.h. to treat a 300 ft. of swath, calibrate equipment to deliver 4.5 gallons per minute.

### ACTIVE INGREDIENTS

Pyrethrins	0.1%
Piperonyl Butoxide, technical	1.0%
Petroleum distillate	0.4%

### INERT INGREDIENTS

98.5%

\*Equivalent to 0.8% (butyl arbutyl (6-propylpiperonyl) ether and 0.2% related compounds.

**CAUTION:** Harmful to fish and aquatic life. Avoid contact with fish and aquatic life. Do not use in or near water. Do not use in or near water. Do not use in or near water. Do not use in or near water. Do not use in or near water.

EPA Reg. No. 1057-64

EPA Reg. No. 1057-64

ACCOUNTED  
1057-64