

# DOLGE GENERAL PURPOSE AQUEOUS PYRETHRUM INSECTICIDE

## DIRECTIONS

### INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated area and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

**SURFACE SPRAYING:** To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and cadellids. Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles. Spray edges of carpeting and under carpeting and rugs. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs. Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat treatment as needed.

To control fleas and brown dog ticks. Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths. Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining. Repeat application at 30-day intervals if treated clothes are not stored in mothproof containers.

### A CONCENTRATED MICRO EMULSION

- CLEAR
- STABLE
- LEAVES NO OILY OR OBJECTIONABLE RESIDUE

EFFECTIVE, FAST ACTING INSECTICIDE.

FOR USE IN HOMES, INSTITUTIONS, RESTAURANTS.

FOOD PROCESSING PLANTS, DAIRIES, LIVESTOCK AND OUTDOORS FOR MOSQUITO CONTROL.

APPLY WITH CONVENTIONAL MECHANICAL OR COMPRESSED AIR EQUIPMENT.

**SPACE SPRAYING AND FOGGING:** Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths, direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 3.6 ounces per 1000 cubic feet of space.

**OUTDOOR ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant ponds, marshy areas, ponds and shore lines may be treated. Application of this product to any body of water is prohibited. Apply at a rate of 1 1/2 gallons per acre (4 1/2 ounces per 1000 square feet).

Using truck-mounted conventional misting equipment such as Biotomistâ models 51C, B7 or 100 HT, traveling at 5 m.p.h. to treat a 100 foot swath, calibrate equipment to deliver 4.5 gallons per minute.

### ACTIVE INGREDIENTS

Pyrethrins

0.1%

\*Piperonyl Butoxide, technical

1.0%

INERT INGREDIENTS

98.9%

\*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

**CAUTION:** Harmful if swallowed. Avoid contamination of food and feedstuffs.

Remove pets, birds and cover fish aquaria before spraying.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on the label.

Do not reuse empty container. Destroy by perforating or crushing and burning in a safe place.

## CAUTION

KEEP OUT OF REACH OF CHILDREN

See Additional Precautions On Right Hand Panel

THE C. E. DOLGE COMPANY

WESTPORT, CONNECTICUT 06897

FPA Reg. No. 1057-34

BEST AVAILABLE COPY

Not Reviewed. Registrant claims to be in accordance with reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.