

Reg # 1057-64

PM-10

File
13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR 6 1994

Barbara P. Minor
UNYTEX RESEARCH CORP.
Agent for: C.B. Dolge Co.
P. O. Box 818
Rochester, NY 14603

OFFICE OF
PREVENTION, PESTICIDES, AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 02/23/94 in Response to PR Notice 93-7
EPA Reg. No. 1057-64
DOLGE GENERAL PURPOSE AQUEOUS [YRETHRUM INSECTICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.

BEST AVAILABLE COPY



Recycled/Recyclable
Printed with Soy/Canola ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

33

Dolge General Purpose Aqueous Pyrethrum Insecticide is a clear, stable, concentrated micro emulsion. It leaves no oily or objectionable residues and is an effective, fast acting insecticide. Dolge General Purpose Aqueous Pyrethrum Insecticide should be applied with conventional mechanical or compressed air equipment.

Dolge General Purpose Aqueous Pyrethrum Insecticide is designed for use in Food Processing Plants, Offices, Restaurants, Bakeries, Warehouses, other Food Establishments, Medical facilities, Office buildings, industrial establishments, Institutions and Schools other than preschool or day care.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets, birds and cover fish aquaria before space spraying or surface applications. Do not directly spray on nursing puppies or kittens. Refer to the Dolge General Purpose Aqueous Pyrethrum Insecticide Material Safety Data Sheet for other safety information.

PERSONAL PROTECTIVE EQUIPMENT:

Applicators and other handlers must wear:
• Long-sleeved shirt and long pants
• Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS: Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, ponds, streams, tidal marshes or estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

EPA Est. No. 527-IL-1

EPA Est. No. 527-NY-1

FORM 1107-B-PROPOSED 100
(REV. 1/21/84)

ACCEPTED
with COMMENTS
In EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
1052-64

BEST AVAILABLE COPY

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply the product in a way that will contact users or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult Mushroom Flies, Fungus Gnats and Cecid Flies in growing houses, apply at a rate of 3-6 ounces per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants, use at a rate of 3-6 ounces to 2000 cubic feet of space.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product which are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not remain in treated areas. Vacate treated areas and ventilate before reoccupying. In medical centers, inform patients prior to treatment. Ventilate rooms for two hours, before returning patients.

INDOOR APPLICATION

IN FOOD PROCESSING ESTABLISHMENTS: The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. For maximum effectiveness, a combination of spot, surface treatment and space spraying is recommended.

(continued on right panel)

DIRECTIONS FOR USE (Continued from left panel)

SURFACE SPRAYING: To control cockroaches, palmetto bugs, ants, silverfish, spiders, crickets, clever mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and cadellids: Use a good sprayer adjusted to a coarse wet spray. Direct spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers, behind and beneath sinks, stoves, refrigerators and cabinets, along baseboards, around door and window frames, into dark corners and in and around garbage compactors, garbage chutes and cans. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

In case of infested foods in cabinets, dispose of the infested material. Remove all contents and clean thoroughly. Spray interior surfaces of empty cabinets thoroughly. After spray has dried, cover shelves with paper or otherwise protect foods from direct contact with sprayed surfaces.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shagging. Repeat treatment as needed.

PET AREAS: To control fleas and brown dog ticks on premises. Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

SPACE SPRAYING: Will kill accessible, exposed stages of flies, including fruit flies, mosquitoes, silverfish, flying moths, gnats, crickets, cockroaches, spiders, scorpions, wasps, hornets, clever mites, cheese mites, cheese skippers and these common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths, meal moths and grain mites. Close doors, windows and transoms and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space. Keep the area closed for at least 15 minutes. Ventilate the area after treatment is complete. Repeat treatment as necessary to aid in the prevention of breeding, spray on standing water and in garbage pails.

DOLGE GENE AQUE PYRETHRUM

EPA Reg. N

ACTIVE INGREDIENTS:

Pyrethrins
• Piperonyl Butoxide, Technical
INERT INGREDIENTS

*Equivalent to 0.8% (butylcarbyl) (8-pr compounds).

KEEP OUT OF REACH CAUTION

STATEMENT OF PRACTICE

If swallowed: Drink one or two glasses of back of the throat with finger. Repeat until vomit not induce vomiting or give anything by mouth. Remove victim to fresh air. Apply artificial respiration if necessary. Remove contaminated clothing and wash off eyes: Flush eyes with plenty of water. Call Emergency Telephone Number: 718

See Left Panel For Additional

THE C.B. DOLGE
P.O. BOX 1515 • ROCHESTER
BRANCH WAREHOUSES

1 U.S. G

NET CONTENTS: 3.8 Lites

ON LIVESTOCK: To protect and gnats, apply a light mist to horses, mares and deer flies, thoroughly. Repeat treatment. To control bloodsucking flies, the spray to the base of the horse. To control poultry lice, spray by spraying over the birds with the legs, under body and tail. For control of bedbugs and other insects in nests where the

FLEAS AND TICKS ON DOGS: eyes, nose, and mouth as you the legs, under body and tail. Make sure spray wets thorough

ADULT MOSQUITO CONTROL: best results, treat when the vegetation where mosquitoes breed. Apply at a rate of 2.5

Using truck-mounted conventional traveling at 5 m.p.h. to treat

OUTDOORS: Camping, tent mosquitoes, midges, gnats, mosquitoes. If in rare cases a good sprayer which will produce

STORAGE: Do not contaminate water, food or feed with PESTICIDE STORAGE. Avoid exposure to extreme material such as sand, sand. **PESTICIDE DISPOSAL:** to label instructions must be CONTAINER DISPOSAL or puncture and dispose of DO NOT

55 Gallons

200 Gallons