

DIRECTIONS FOR USE:

MIXING. CROPS: Use Valley Brand Malathion Emulsifiable No. 5 at the rate recommended below. For maximum control, apply at intervals of 5 to 7 days, or as necessary to keep insects under control. The rate and interval of application will depend upon the insect being controlled, the type of infestation, weather, and method of application. Proper timing and method of application are essential for best results.

USE ON CROPS: Use Valley Brand Malathion Emulsifiable No. 5 at the rate recommended below. For maximum control, apply at intervals of 5 to 7 days, or as necessary to keep insects under control. The rate and interval of application will depend upon the insect being controlled, the type of infestation, weather, and method of application. Proper timing and method of application are essential for best results.

COTTON:

BOLL WEEVIL AND GRASSHOPPERS: Apply at rate of one (1) gallon per 2 1/2 to 5 acres.

THRIPS:

Apply at rate of one (1) gallon per 5 to 16 acres.

COTTON APHIDS, LYCUS BUGS, TARNISHED PLANT BUGS: —

Apply at rate of one (1) gallon per 2 to 3 acres.

COTTON FLEAHOPPERS:

Apply at rate of one (1) gallon per 4 to 7 acres.

GARDEN WERWORMS:

Apply at rate of one (1) gallon per 2 1/2 to 7 acres.

VEGETABLES:

BEANS:

Apply at rate of one (1) gallon per 4 acres.

CUCUMBERS:

Apply at rate of one (1) gallon per 4 acres.

TOMATOES:

Apply at rate of one (1) gallon per 4 acres.

POTATOES:

Apply at rate of one (1) gallon per 4 acres.

PEAS:

Apply at rate of one (1) gallon per 5 to 10 acres.

PEAS:

Apply at rate of one (1) gallon per 4 to 5 acres.

WHEN MAKING APPLICATIONS WITH AIRPLANES: —

Apply at rate of one (1) gallon per 2 to 3 acres.

RESTRICTIONS:

VEGETABLE CROPS: —

Apply at rate of one (1) gallon per 4 acres.

PEAS:

Apply at rate of one (1) gallon per 5 to 10 acres.

PEAS:

Apply at rate of one (1) gallon per 4 to 5 acres.

PEAS:

Apply at rate of one (1) gallon per 5 to 10 acres.

PEAS:

Apply at rate of one (1) gallon per 4 to 5 acres.

PEAS:

Apply at rate of one (1) gallon per 5 to 10 acres.

PEAS:

Apply at rate of one (1) gallon per 4 to 5 acres.

PEAS:

Apply at rate of one (1) gallon per 5 to 10 acres.

PEAS:

Apply at rate of one (1) gallon per 4 to 5 acres.

PEAS:

Apply at rate of one (1) gallon per 5 to 10 acres.

PEAS:

Apply at rate of one (1) gallon per 4 to 5 acres.

PEAS:

Apply at rate of one (1) gallon per 5 to 10 acres.

MALATHION EMULSIFIABLE No. 5

(CONTAINS 5 POUNDS MALATHION PER GALLON)

ACTIVE INGREDIENTS:

Malathion	5.00%
Acesulfone	52.00%
INERTS	43.00%
TOTAL	100.00%

**O.O-Dimethyl

phosphorothioic acid, dimethyl ester, succinate

HARMFUL TO

MAN, DOMESTIC ANIMALS, AND PLANTS

DO NOT USE THIS PRODUCT FOR ANY PURPOSES

OTHER THAN THOSE SPECIFIED ON THE LABEL

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

BACK RUBBING DEVICES:

Use 2 to 3 gallons of Malathion Emulsifiable No. 5 per acre. For maximum control, apply at intervals of 5 to 7 days, or as necessary to keep insects under control. The rate and interval of application will depend upon the insect being controlled, the type of infestation, weather, and method of application. Proper timing and method of application are essential for best results.

HOGS:

Apply at rate of one (1) gallon per 5 to 10 acres.

RESTRICTION ON LIVESTOCK:

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

DO NOT FEED TO ANY ANIMALS FOR 24 HOURS AFTER APPLICATION.

8-24-67

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

1063-68

CALL A PHYSICIAN

IMMEDIATELY

KEEP OUT OF REACH OF CHILDREN

SEE OTHER WARNING ON RIGHT PANEL

VALLEY CHEMICAL COMPANY

GREENVILLE, SOUTH CAROLINA

Manufactured by

VALLEY CHEMICAL COMPANY

GREENVILLE, SOUTH CAROLINA

DO NOT USE THIS PRODUCT FOR ANY PURPOSES

OTHER THAN THOSE SPECIFIED ON THE LABEL

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

EXTERNAL: WASH SKIN WITH SOAP AND WATER

INTERNAL: DRINK MILK OR EGGS

ANTIDOTE: INGESTION OF WARM WATER, INDUCE VOMITING

THE UNIVERSITY OF MICHIGAN
LIBRARY

GREENGLASS P.O. BOX 30 MILWAUKEE

NOTICE TO USER: This document contains information that is exempt from public release under the Freedom of Information Act, 5 U.S.C. 552. It is being furnished to you for your information only. It is not to be distributed outside your agency. If you are not the intended recipient, please do not disseminate, copy, or otherwise use this information. If you have received this document in error, please notify the person or agency to whom the document was addressed.