NIROLLED.

SHELLED OR EARL BARLEY , SORGHUM, WHEAT AND RYE: I FILES and MOTHS: FOR GRAIN A LOES INTO STORAGE, HAND BOTH IN The state of the s The State of the S FOR

G. A. ALREADY IN STORAGE, maxing and the contract of the first of the contract Commence of the second of the The state of the s DO NOT ...

> THE BEETLES and MOTHS: FOR GRAIN "TO STORAGE, " The state of the

FOR JRAIN TORAGE.

TAS FILCORN BARLLY OATS, SORGHUM THE ROLL and COTTON SEEDS BEETLES on a

OF APPLANTAGE OF PERTIAS and MOTHS



Malathion Grain Protectant

(Contains 5 pounds Premium Grade Malathion per Gallon)

ACTIVE INGREDIE	ints:	BY WEIGHT	
Malathion**	 	₩.75°	
Arometic Petrol	eum Solvent	37.5 ?	
TOTAL			
-Dimethyl Di	thiophosphate of Dieth	yl Mercaptosucc	
	Y SWALLOWING, OR SKIN CONTACT		
wate	NAL: — Give a tabl r; repeat until vomit d keep quiet.	espoonfu! fluid is	
	skin is contacted, raiately wash with soa		
CALLA	PHYSICIAN IMM	EDIATI :	
	CAUTION		

KEEP OUT OF REACH OF CHILDREN

THE OTHER WARNINGS ON RICHT PANEL

--ctu-

VALLEY CHEMICAL COMPANY

THE AN THE STORT OF A PROPERTY.

13413 1.

Beetle, Rice Weevil, Granary Weevil, Sawtoothed Grain Beetle, Flat Grain Beetle, Red Flour Beetle, Rusty Grain Beetle, Lesser Grain Borer, and Indian Meal Moth.

CORN (SHELLED OR EAR), BARLEY, OATS, SORGHUM, WHEAT AND RYE:
BEETLES and MOTHS: — FOR GRAIN AS GOES INTO STORAGE, mix 1 pint of Malathion Grain Protectant in 2 to 5 gallons of water to treat 1,000 bushels. Spray into grain stream as it goes into storage. FOR GRAIN ALREADY IN STORAGE, mix 12 pint Malathion Grain Protectant in 1 to 2 gallons of water to treat 1,000 sq. ft. of grain surface. Will not control insects established beneath the surface. DO NOT repeat surface treatment.


RICE (ROUGH): — BEETLES and MOTHS: — FOR GRAIN AS GOES INTO STORAGE, mix 1 pint of Malathion Grain Protectant in 2 to 5 gallons of water to treat 1,000 bushels. Spray into grain stream as it goes into storage. FOR GRAIN ALREADY IN STORAGE, mix 12 pint of Malathion Grain Protectant in 1 to 2 gallons of water to treat 1,000 sq. ft. of grain surface. Will not control insects established beneath the sarface. DO NOT repeat surface treatment


STORAGE BINS:—For CORN, BARLEY, OATS, SORGHUM, WHEAT, RYE, RICE and COTTON SEED:—BEETLES and MOTHS: Treat storage bins wall and floor 2 to 4 weeks before grain is put into bin. Mix 10 ozs. Malathion Grain Frotestant in 2 gallons of water and spray 1,000 sq. feet.

FIELD and GARDEN SEEDS: — BEETLES and MOTHS: I will and garden seeds can be protected against these insects with a dosage of 1 pint Malathion Grain Protectant in 1 to 2 willons of water and treating 500 bushels of seed. Do not use tood or feed after treating at this rate.

J. S.D.A. Reg. No. 1063-113







THIS PRODUCT FORMUL PREMIUM GRADE MALA! NICAL.

CAUTION: riamintul is swal or absorbed arough the skining spray mist. Avoid cont eyes or eathing. Wash thorou and water after using. Avoid of feed and focastuffs, food watering troughs. Keep out children and domestic anin Grain Protectant upon reper or careless use, may cause ch hibition. If poisoning occur medical aid. Atropine is an use where milk is processed move lactating animals from treating. Also remove calv month of age before treating

CAUTION: — Malathion Grain Protectant in certain species of fish. Avoid contamination and streams.

CLEAN UP SPILLAGE AND SPRAY WAS: liquid with sawdust or other absorbent mate Clean up area with strong lye solution.

DISPOSAL OF EMPTY CONTAINERS:

completely wash the outside of the empty water. Rinse out the inside of the container which has been added household or commercial caustic soda (lye). Carefully discard the rindurying at least 18 inches deep in an isolated water supplies. Fifty-five gallon drums may toof to any approved drum reconditioner: sm should be broken, mutilated and punctured, a may be buried at least 18 and es deep in a away from water supplies.

NOTICE TO USERS. Selier worrant that the to its chemical description and is reasonably fit for ed on the label when used in accordance with direct conditions of u.e. but neither the warranty nor are of merchantability or fitness for a particular purpoplied, extends to the use of this product contrary to or under abnormal conditions, or under condition foreseeable to eller, and boxer as are the release of

Confused Flour y Weevil, Sawain Beetle, Red Beetle, Lesser al Moth.


AR), BARLEY, T AND RYE:


— FOR GRAIN, mix 1 pint of in 2 to 5 gallons hels. Spray into storage. FOR RAGE, mix 12 ctant in 1 to 2 0 sq. ft. of grain sects established T repeat surface

it goes into storage. FOR GRAIN mix 1 pint of Malathion Grain of water to treat 1,000 bushels. It goes into storage. FOR GRAIN mix 12 pint of Malathion Grain of water to treat 1,000 sq. ft. of tol insects established beneath the face treatment.

th. Barley, Oats, sorghum, COTTON SEED:—BEETLES and bins wall and floor 2 to 4 weeks in. Mix 10 ozs. Malathion Grain water and spray 1,000 sq. feet.

DS: — BEETLES and MOTHS: — be protected against these insects lathion Grain Protectant in 1 to 2 ag 500 bushels of seed. Do not use g at this rate.





INER

O,O

inate

HAR

ANTIDOTE:

glass of warm
patient lie down

EXTERNAL: —
clothing and imme

in the contaminated and water.

SEE OTHER WARN ON RIGHT PANEL

Manufa ared by

VALLEY C

CALL A

CHEMICAI

COMPANY

DIATELY

REENVILLE

P O BOX 1317

MISSISSIPPI

THIS PRODUCT FORMULATED WITH PREMIUM GRADE MALATHION TECHNICAL.

CAUTION: Harmful is swallowed, inhaled or absorbed chrouge the skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed and foodstuffs, food containers and watering troughs. Keep out of the reach of children and domestic animals. Malathion Grain Protectant apon repeated, prolonged or careless use, may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal. Do not use where milk is processed and stored. Remove lactating animals from building before treating. Also remove calves under one month of age before treating.

CAUTION: — Malathion Grain Protectant may be toxic to certain species of fish. Avoid contamination of lakes, ponds and streams.

CLEAN UP SPILLAGE AND SPRAY WASTE: — Soak up liquid with sawdust or other absorbent materials and bury. Clean up area with strong lye solution.

DISPOSAL OF EMPTY CONTAINERS: - Carefully and completely wash the outside of the empty container with water. Rinse out the inside of the container with water, to which has been added household or commercial detergent and caustic soda (lye). Carefully discard the rinse solution by burying at least 18 inches deep in an isolated area away from water supplies. Fifty-five gallon drums may then be disposed of to any approved drum reconditioner; smaller containers should be broken, mutilated and punctured, after which they may be buried at least 18 inches deep in an isolated area away from water supplies.

NOTICE TO USERS: - Selier warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or timess for a particular purpose, express or insplied, extends to the use of this product contrary to label instruction or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the rick of such use