

[1074-18]

Kills advanced stages of granary weevil, rice weevil, cadelle, corn moth, Indian meal moth, saw-toothed grain beetle, angoumois grain meal moth and certain other common grain insects.

- Terminals • Elevators • Warehouses
- Farm Storage • Bunkers • Seed Stores
- Flat Storage • Mills • Food Stores

Can be used in food processing plants, bakeries, creameries, dairies, hotels and homes.

### DIRECTIONS

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY

**SOUTHERN MILL SPRAY** is an efficient insecticide for use in households, grain elevators, flour mills and food processing plants. When properly used **SOUTHERN MILL SPRAY** will control grain moths, grain weevils, grain mites, silverfish, roaches, and certain other insects that infest food processing establishments. Before spraying thoroughly clean areas to be treated. Applications should be made to exposed stages of insects to insure best results.

**SURFACE SPRAY:** Treat walls, floors, ceilings, conveyors and other surfaces with Southern Mill Spray at the rate of 1 gallon per 1000 square feet. Cracks and crevices should be treated heavily and thoroughly.

**GRAIN TREATMENT:** **SOUTHERN MILL SPRAY** may be applied directly to the surface of grain. Grain should be leveled and application made after fumigation to prevent moths and other insects from infesting grain surface causing subsequent caking and wetting. Apply 1 gallon of **SOUTHERN MILL SPRAY** per 750 square feet of grain surface. Repeat as required for continued control.

NET CONTENTS

GALLONS

# SC-PRO-CO

# SOUTHERN MILL SPRAY

ACTIVE INGREDIENTS 100%

Pyrethrins	0.15%
Technical piperonyl butoxide*	1.50%
Petroleum distillate	98.35%

\*Consists of 1.2% (butyl carbityl) (6-propylpiperonyl) ether and 0.3% related compounds

### CAUTION

KEEP OUT OF THE REACH OF CHILDREN  
See caution statement side panel

## SOUTHERN PRODUCTS COMPANY

1307 Eastman Pl. S. 6th & Davidson St.  
Memphis, Tenn. Nashville, Tenn.

### FOR INDUSTRIAL USE

**SPACE SPRAY:** To control flies, gnats, mosquitoes and other flying insects, close doors and windows before spraying. Apply 1 gallon of Southern Mill Spray per 80,000 cubic feet of space, allowing doors to remain closed for at least 15 to 20 minutes after treatment. Repeat as required for continued control.

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product as a space spray in these areas, apply only when plant is not in operation. Do not remain in treated areas. Ventilate the area after treatment is completed.

### CAUTION

Do not apply directly to food products. Avoid inhalation of spray mist, contact with skin or eyes. Harmful if swallowed. Keep away from children. Mixture as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place. Do not use, pour, spill or store near heat or open flame.

### NOTICE

The above directions are based upon use under normal and reasonably foreseeable conditions. If the directions are followed and precautions on this label observed, available research indicates that this product will give satisfactory results. However, because climatic, geographic, and other conditions of use will vary widely, the seller makes no warranty or representation, expressed or implied, concerning this product or its use except that it conforms to the chemical description on the label and no person is authorized to make any other representations upon their behalf. The seller shall not be held responsible for damages resulting from the handling, storage, or use of this product for any reason except the failure of the material to conform to the chemical description on this label. Damages, if any, which may result from the failure of this product to conform to the description on this label shall be limited to direct damages but shall not include consequential damages such as those resulting from crop injury, or failure to give adequate control.

U.S.D.A. REG. NO. 1074-18

ACCEPTED

Aug. 30, 1971

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROUQUENTICIDE ACT  
FOR ECONOMIC PESTICIDE REGISTERED  
UNDER 190

1074-18

[1074-18]

**Kills exposed stages of granary weevil, rice weevil, cadelle, conmoth, Indian fusea flour beetle, saw toothed grain beetle, angou.nois grain meal moth and certain other common grain insects.**

- Terminals      • Elevators   • Warehouses
- Farm Storage   • Boxcars   • Seed Stores
- Flat Storage   • Mills       • Feed Stores

Can be used in food processing plants, bakeries, creameries, dairies, hotels and homes.

**DIRECTIONS**

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY

SOUTHERN MILL SPRAY is an efficient insecticide for use in households, grain elevators, flour mills and food processing plants. When properly used SOUTHERN MILL SPRAY will control grain moths, grain weevils, grain mites, silverfish, roaches, and certain other insects that infest food processing establishments. Before spraying thoroughly clean areas to be treated. Applications should be made to exposed stages of insects to insure best results.

**SURFACE SPRAY:** Treat walls, floors, ceilings, conveyors and other surfaces with Southern Mill Spray at the rate of 1 gallon per 1000 square feet. Cracks and crevices should be treated heavily and thoroughly.

**GRAIN TREATMENT:** SOUTHERN MILL SPRAY may be applied directly to the surface of grain. Grain should be leveled and application made after fumigation to prevent moths and other insects from infesting grain surface causing subsequent caking and wetting. Apply 1 gallon of SOUTHERN MILL SPRAY per 750 square feet of grain surface. Repeat as required for continued control.

**FOR INDUSTRIAL USE**

**NET CONTENTS**

**GALLONS**

# SO-PRO-CO SOUTHERN MILL SPRAY

<b>ACTIVE INGREDIENTS</b>	<b>100%</b>
Pyrethrin	0.15%
Technical piperonyl butoxide	1.50%
Petroleum distillate	98.35%
Consists of 1.2% (butylparbityl) (6-propylpiperonyl) ether and 0.3% related compounds.	

**CAUTION**  
KEEP OUT OF THE REACH OF CHILDREN  
See caution statement side panel

**SOUTHERN PRODUCTS COMPANY**  
1301 Heistan Pl.      S. 6th & Davidson St.  
Memphis, Tenn.      Nashville, Tenn.

ACCEPTED

*[Handwritten signature]*

U.S. DEPT. OF AGRICULTURE

**SPACE SPRAY:** To control flies, gnats, mosquitoes and other flying insects, close doors and windows before spraying. Apply 1 gallon of Southern Mill Spray per 50,000 cubic feet of space, allowing doors to remain closed for at least 15 to 20 minutes after treatment. Repeat as required for continued control.

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product as a space spray in these areas, apply only when plant is not in operation. Do not remain in treated areas. Ventilate the area after treatment is completed.

**CAUTION**

Do not apply directly to food products. Avoid inhalation of spray mist, contact with skin or eyes. Harmful if swallowed. Keep away from children. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place. Do not use, pour, spill or store near heat or open flame.

**NOTICE**

The above directions are based upon use under normal and reasonably foreseeable conditions. If the directions are followed and precautions on this label observed available research indicates that this product will give satisfactory results. However, because climatic, geographic, and other conditions of use will vary widely the seller makes no warranty or representation, expressed or implied, concerning this product or its use except that it conforms to the chemical description on the label and no person is authorized to make any other representations upon their behalf. The seller shall not be held responsible for damages resulting from the handling, storage, or use of this product for any reason except the failure of the material to conform to the chemical description on this label. Damages, claims, or charges, resulting from the failure of this product to conform to the description on this label shall be limited to direct damages but shall not include consequential damages, such as those resulting from crop loss or the failure to obtain adequate control.

U.S.D.A. REG. NO. 1074-18

*[Faint handwritten text]*