

DIRECTIONS

For best results use Weed Spray when the weeds are young and in a rapid growing condition. For perennial weeds and under conditions where weeds are difficult to kill use the higher rates.

WEEDS, SPRING AND WINTER: Apply Weed Spray at the rate of 1 quart to 2 quarts per acre in 100 gallons of water as soon as weeds appear.

WEEK FLAME CONTROL: For certain 2,4-D susceptible species such as Alder, Aspen, Cat-barked, Buckbrush, Elderberry, Hazel, Locust, Sassa, Wild Grape and Willow. Apply at a rate of 2 quarts in 100 gallons of water as a foliar application, using a drenching spray soon after leaves are fully developed. Complete coverage is important and all parts of the plant should be sprayed to the point of run-off. Do not treat during period of severe drought or in early fall when leaves have lost their healthy color.

TENN AND GOLF COURSES: Apply 2 pints of Weed Spray per acre in sufficient water to provide uniform coverage. Do not apply on creeping grasses such as bent or to newly seeded turf areas until grasses have become well established. Against Wild Garlic and Wild Onion apply 2 quarts to 3 quarts per acre in very early spring. Repeat treatments will be necessary annually for at least three years.

NOTICE: We guarantee the contents of this package to conform to the labeled composition. The use of this material being beyond our knowledge and control, we do not make any warranty, express or implied, as to the effects of such use whether or not in accordance with the directions.

NET CONTENTS

20-750-0
SOUTHERN
WEED SPRAY
Contains 2,4-D
4 lbs. per Gallon

GALLONS

ACCEPTED
MAY 7 1969.
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTE-
ED UNDER NO. 1074-19 SUBJECT
TO ATTACHED COMMENTS.

Low Volatile Emulsifiable Concentrate

ACTIVE INGREDIENTS

2-Ethylhexyl Ester of 2,4-Dichlorophenoxyacetic acid (Equivalent to 2,4-Dichlorophenoxyacetic acid 4.4%)..... 67.0 %

INERT INGREDIENTS:..... 33.0 %
100.0 %

CAUTION
Keep Out of Reach of Children
May Be Harmful If Swallowed
See Caution On Side Panel

U.S.D.A. REG NO 1074-19

SOUTHERN PRODUCTS COMPANY
MEMPHIS, TENN NASHVILLE, TENN.

This product is recommended for control of Arrowweed, Sandbolic, Dock, Bindweed, Nutsedge, Lambsquarters, Nettle, Mustard, Purslane, Pigweed, Plantain, Ragweed, Shepherd's-purse, Thistle, Wild Garlic, Wild Onion-Glory, Wild Garlic, Wild Onion, and many other broad leaved weeds on golf courses, in parks and other recreational areas, along highways, on airfields, on vacant lots and other waste areas where broad leaved weed control is desired.

Fill spray tank with half the required quantity of water, then add the recommended amount of Weed Spray with agitation and continue filling the spray tank with water. The amount of water required in low volume applications may vary from 5 to 25 gallons per acre. For high volume applications, 50 to 100 gallons of water or more may be necessary for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

CAUTION: Avoid spray drift to susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees, and ornamentals. Spray drift may cause severe injury to both growing and dormant plants. Coarse sprays are less likely to drift. Although this product is formulated with low volatile esters, high temperatures may produce vapors after application which might injure 2,4-D sensitive plants growing nearby.

Do not contaminate irrigation ditches or water used for domestic purposes. Do not use the same spray equipment for applying other materials to plants. Do not store near fertilizers, seed, insecticides, or fungicides. Do not take internally. Avoid contact with skin, eyes or clothing. Harmful if swallowed.

Cameraman's Note
Evan Gray