## USE DIRECTIONS

For Household and Hospital use on floors, walls and woodwork.

Remove all gross filth, dirt and heavy soils prior to application of the disinfectant.

Wet surfaces thoroughly with disinfectant.

#### SANITIZING SOLUTION

One ounce SG 250 to one gallon of water.

#### DISINFECTING SOLUTION

For household, broad spectrum disinfecting use 3 Oz. SG 250 to  $1\frac{1}{4}$ gallons water.

#### HOSPITALS

Use 31/2 ounces to 11/4 gallons of water as hospital disinfectant. Used as directed will be effective against pseudomonas aeruginosa PRD 10-

## FOOD PROCESSING EQUIPMENT

Clean and rinse equipment thoroughly, then apply sanitizing solution. One ounce SG 250 per 1 gallon of water. No rinse.

### DAIRIES

Clean and rinse equipment thoroughly, then apply sanitizing solution containing 1 oz. SG 250 to 1 gallon of waler (195 ppm). No rinse required. Follow recommendations of Local Health Board.

## NET CONTENTS

SO-PRO-CO

# SG 250 CONCENTRATE

Dilute Before Using

**ACTIVE INGREDIENTS:** 

n—Alkyl (60%  $C_{14}$ , 30%  $C_{16}$ , 59 5% C<sub>18</sub>) dimethyl benzyl ammoni n-A(68% C<sub>12</sub>, 32% C<sub>14</sub>) dimethy HK/ ammonium chlorides INERT INGREDIENTS

WARNIN

Keep out of reach of ch See other warning on side panel.

EPA Reg. No. 1074-34

CONTRACTOR DOCOLICTC SOUTHERN PRODUCTS COMPANY 1301 Heistan Pl. Memphis, Tenn.


## GALLONS





% C <sub>12</sub> , iium chlorides yl_ethylbenzyl	
	1.25%
····· · · · · · · · · · · · · · · · ·	97.50%
	100.00%
G nildren.	





## **RESTAURANT AND BAR**

Rinse Dishes, glassware, Silverware, Cooking Utensils, Wash with soap or synthetic detergent, rinse thoroughy and immerse in a solution containing 1 oz. SG 250 to 1 gallon of water (195 ppm)

RINSE EMPTY CONTAINER THO-ROUGHLY WITH WATER AND DIS-CARD.

WARNING ---- Keep out of reach of Children. Do not get in eyes, skin, or on clothing. Causes severe eye irritation. Causes skin irritation. In case of contact, wash immediately with plenty of water. For eyes get medical attention.

## FIRST AID

If swallowed, do not induce vomit ing. Drink large quantities of fluid and call a physician immediately.