

[1075-6]
2 pages

ALEXO

FOGGING INSECTICIDE

For Use on: Flies, Mosquitoes, Beetles, Fleas, Lice, and other household pests, and on
Barnacles and Food Processors and Equipment.

Apply with Equipment Designed for use with Fogging Insecticides. Do Not
Expose to Heat, Acids, Alkalis, or Oxidizing Agents.

The insecticide component of ALEXO fogging is classified as a fast-acting contact insecticide
as directed, when used in accordance with the following instructions.

| | |
|--------------------------------|------|
| ACTIVE INGREDIENTS: | 100 |
| Piperonyl Butoxide | 0.3 |
| *Piperonyl Butoxide, Technical | 0.0 |
| Petroleum Distillate | 99.7 |

*Equivalent to 2.4-dichlorobiphenyl 0.6 propylparathion 0.001 and 0.6 of related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Keep far from food, feed, and feed stuffs. Avoid contamination of feed and feed stuffs.
Remove pets, birds, and other farm animals before spraying.
The product is toxic to fish. Keep out of rivers, streams, or ponds. Do not contaminate
water by cleaning of equipment or disposal of wastes. Apply this product only as
specified on this label.
Do not reuse empty drum. Return to manufacturer or distributor, by perforating or
crushing and burying in a safe place away from water supplies.
Do not store near open flame. Do not use in garages, or similar areas.
Do not use in the vicinity of homes where infants, children, and persons are confined.
Do not breathe vapors. Use only in the manner of application recommended by the
U.S. Bureau of Entomology and Plant Quarantine and the material. Wash hands,
arms, and face with soap and water after handling and before eating or smoking. Wash
all contaminated clothing with soap and hot water before reuse.

Net Contents: Gallons

THIS PRODUCT MUST BE USED ONLY BY TRAINED PERSONNEL !!!

© 1975, 10, 1075-6
EPA EST. NO. 1075-NY-01



MANUFACTURED BY
ALEXANDER CHEMICAL CO.
36 WALWORTH STREET, BROOKLYN 3, NEW YORK

AUG 8 1975
1075-4

[1075-6]

DIRECTIONS FOR USE

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

SPACE SPRAYING:

Close windows and doors and shut off ventilating systems. Use a Fogger or a generator adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of flies, fruit flies, mosquitoes, small flying moths, gnats, wasps, hornets, and these common stored product pests: Angoumois grain moths, Mediterranean flour moths, Indian meal moths, Tobacco moths, and meal moths.

NOTE: WASPS AND HORNETS application should be made in late evening when insects are at rest. Spray into hiding and breeding places contacting as many insects as possible.

SURFACE SPRAYING:

To control exposed stages of roaches, ants, silverfish, virus, rice weevils, confused flour beetles, rust red flour beetles, saw toothed, grain beetles, spider beetles, cigarette beetle, Drugstore beetles, meal worms, grain mites, and cadellies: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silver fish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages, directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

To control carpet beetles: Spray edges of carpeting and under carpeting and mats. Make localized application to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

Whenever possible directly contact the pests with the spray. In commercial and industrial type applications, infested food products should be fumigated or treated by an approved product.

In the home, infested food should be disposed of completely. Shelves and drawers cleaned, sprayed so as to contact all pests. Then covered with shell paper after allowing to dry. Utensils contacted by spray should be washed before using.

THIS PRODUCT MUST BE USED ONLY BY TRAINED PERSONNEL