

DIRECTIONS FOR USE

GALLONS



U. S. PATENT PENDING

ACTIVE INGREDIENTS

Copper salts of fatty acid esters
 6.00 percent by weight

INERT INGREDIENTS

94.00

100

100

100

CITIES SERVICE COMPANY

AMERICAN COPPERHILL OPERATIONS
COPPERHILL, TENNESSEE

FRUIT TREES
 Apply to all fruit trees for control of scale, aphids, and other sucking insects. Apply in early spring and repeat as needed.

WALNUTS - Blight
 Apply to all walnuts for control of blight. Apply in early spring and repeat as needed.

VEGETABLES

CARROTS - Leaf spot
 Apply to all carrots for control of leaf spot. Apply in early spring and repeat as needed.

CELERY - Early blight
 Apply to all celery for control of early blight. Apply in early spring and repeat as needed.

CUCURBITS (CUCUMBERS, CANTALOUPE OR MUSKMELONS, SQUASH, PUMPKINS, WATERMELONS) - Powdery mildew, Alternaria blight
 Apply to all cucurbits for control of powdery mildew and Alternaria blight. Apply in early spring and repeat as needed.

CUCUMBERS - Downy mildew
 Apply to all cucumbers for control of downy mildew. Apply in early spring and repeat as needed.

LETTUCE - Downy mildew
 Apply to all lettuce for control of downy mildew. Apply in early spring and repeat as needed.

ONIONS - Downy mildew
 Apply to all onions for control of downy mildew. Apply in early spring and repeat as needed.

PEPPERS - Bacterial leaf spot
 Apply to all peppers for control of bacterial leaf spot. Apply in early spring and repeat as needed.

TOMATOES - Early blight, Septoria leaf spot
 Apply to all tomatoes for control of early blight and Septoria leaf spot. Apply in early spring and repeat as needed.

FIELD CROPS

BARLEY - Leaf blight
 Apply to all barley for control of leaf blight. Apply in early spring and repeat as needed.

RED AND NAVY BEANS - Bacterial blight
 Apply to all red and navy beans for control of bacterial blight. Apply in early spring and repeat as needed.

PEANUTS - Leaf spot
 Apply to all peanuts for control of leaf spot. Apply in early spring and repeat as needed.

SOYABEANS - Leaf spot
 Apply to all soybeans for control of leaf spot. Apply in early spring and repeat as needed.

WHEAT - Leaf blight
 Apply to all wheat for control of leaf blight. Apply in early spring and repeat as needed.

HEADEN

HEADEN - Head blight
 Apply to all headen for control of head blight. Apply in early spring and repeat as needed.

HEADEN - Head blight
 Apply to all headen for control of head blight. Apply in early spring and repeat as needed.