


ACCEPTED
 Dec 18, 72
 RECEIVED
 EST. 1900

hockwald



 A CONSOLIDATED FOODS COMPANY • RESPONSIVE TO CONSUMER NEEDS

DIRECTIONS

GENERAL: Prior to sanitation or disinfection, clean all surfaces with a suitable detergent and rinse thoroughly with potable water. Consult your Oxford Technical Sales Engineer regarding the specific detergent needed for each particular application. Follow the recommendation of your local Health Department regarding sanitation and disinfection of surfaces. For use on hard surfaces, the recommended use dilution solution may be applied by wiping, scrubbing or mopping, making sure all surfaces are thoroughly wet.

DISINFECTING SOLUTION: In Hospitals, public rooms, and homes, clean surfaces such as floors, woodwork, wash basins, urinals and toilets with detergent and rinse thoroughly. Use 3½ oz. HYCO-10 to 5 gallons of water. This solution will be effective against *Pseudomonas aeruginosa* PRD-10.

SANITATION OF MEAT PACKAGING AND FOOD PROCESSING EQUIPMENT: Clean and rinse equipment thoroughly, then apply sanitizing solution of 1 oz. HYCO-10 per 4 gallons of water. No rinse required.

NET CONTENTS

HYCO - 10

ACTIVE INGREDIENTS 10%
 n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides . . . 5%
 n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides . . . 5%
 INERT INGREDIENTS 90%
 100%

HYCO-10 is a concentrate and should be diluted before using.

EPA Reg. No. 1111-146-AA

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and severe skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food. **FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantity of water. Avoid alcohol. Call a physician immediately. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

This product is toxic to fish, wildlife and birds. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

LS - 1A

K 72

DIRECTIONS

SANITATION OF DAIRY EQUIPMENT: Clean and rinse equipment thoroughly, then apply sanitizing solution of 1 oz. HYCO-10 per 4 gallons of water. No rinse required.

RESTAURANT AND BAR RINSE: Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent, rinse thoroughly and then immerse in a solution containing 1 oz. HYCO-10 to 4 gallons of water (195 PPM).

SANITATION OF FABRICS: Agitate in a solution containing 1 oz. concentrate per gallon of water to disinfect linen and clothing. To render diapers and other linens bacteriostatic and to retard development of ammonia, immerse in solution of 1 oz. concentrate per 10 pounds of cloths.

SANITATION OF BEAUTY PARLOR AND BARBER SHOP EQUIPMENT: Soak combs, brushes, razors, etc., for 10 minutes in solution of 1 oz. concentrate per gallon of water.

NET CONTENTS _____ GALLONS

MANUFACTURED BY

HOCKWALD CHEMICALS
 A Division of
OXFORD CHEMICALS
 P.O. Box 227, Brisbane, Calif. 94005
 7201 Paramount Blvd., Pico Rivera 90660

ESTABLISHED SINCE 1900