


# hockwald



 A CONSOLIDATED FOODS COMPANY • RESPONSIVE TO CONSUMER NEEDS

Hockwald 1111 is recommended for use in fogging operations to temporarily reduce airborne bacteria. Fogging with 2 ounces of Hockwald 1111 to 1 gallon of water 6 minutes per 1,000 cubic feet of space will effectively reduce bacteria. Allow a minimum of two hours after fogging has stopped before room can be occupied.

A concentrated one-step cleaner and powerful disinfectant-deodorant of proven effectiveness. The combination of multiple germicides and high active ingredients provides an effective one-step method of disinfecting and cleaning at a dilution of 1 ounce to 1 gallon of water (1:128). FOR USE IN SANITATION IN MEAT AND POULTRY PRODUCTION - FOOD PROCESSING - HOSPITALS - ANIMAL AND POULTRY CARE - VETERINARY. VIRICIDAL - SALMONELLACIDAL - STAPHYLOCIDAL - TUBERCULOCIDAL - FUNGICIDAL - PSEUDOMONACIDAL - CONDUCTIVE.

# HOCKWALD ACCEPTED

# 1111

NOV 12 1974

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 1111-149

## BACTERICIDAL DETERGENT

**ACTIVE INGREDIENTS:** Isopropanol, 12.00%; Potassium laurate, myristate and related compounds, 11.80%; Para-tertiary-butylphenol, 3.45%; ortho-phenylphenol, 3.45%; 4-Chloro-2-cyclopentylphenol, 1.87%; Tetrasodium ethylenediaminetetraacetate, 1.0%; Essential oils, 0.40%.

**INERT INGREDIENTS: 66.03%.**

## DANGER

### KEEP OUT OF REACH OF CHILDREN

See side panels for Precautions and Instructions

EPA Reg. No. 1111-149

EPA Est. 1111-CA-1

PHD-6N

NET CONTENTS

GALLONS

J 74

### DIRECTIONS

1. For disinfecting and cleaning in meat and poultry production, such as in processing of meat, poultry, turkeys, and equipment, dilute Hockwald 1111 1 ounce to 1 gallon of water (1:128). Wipe, mop or swab all surfaces with solution. Leave in contact 10 minutes. All food contact surfaces must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.
2. For non-food disinfection of laying and broiler houses and hatcheries: Do not use in milking stalls, milking parlors or milk houses. Remove all animals, poultry and feeds from the premises. Remove all litter and manure from floors and other surfaces. Empty all feeding and watering troughs and appliances. Saturate all surfaces with a 1:128 solution. Ventilate buildings and other closed spaces. Do not house poultry, livestock or employ equipment until treatment has been absorbed or dried. All treated feed racks, waterers and fountains must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.
3. For general cleanup in food processing plants, patient rooms, surgery, nursing homes, all animal facilities, on walls, floors, equipment, use 1 ounce Hockwald 1111 per 1 gallon of water (1:128). Cover surfaces thoroughly with solution and allow to remain in contact 10 minutes. For porous surfaces, increase concentration to 2 ounces per gallon. Rinse all surfaces coming in contact with food with potable water before reuse.
4. For disinfecting and deodorizing garbage and trash receptacles, carts and sick room equipment, use 1 ounce Hockwald 1111 to 1 gallon of water (1 part to 128 of water).
5. Hockwald 1111 is a prophylactic agent in controlling transmission of athlete's foot fungi on environmental surfaces. Prepare a solution of 1 ounce Hockwald 1111 per 1 gallon of water; mop or swab this solution onto floors and other surfaces normally contacted by the feet in locker rooms, shower stalls, bathrooms and around swimming pools.

AC  
Sta  
Sal  
Pse  
AC  
Tri  
\*H  
(st  
virt  
Ho  
gar  
par  
AC  
Sal  
Sal  
Sal  
Sal  
Ho  
gar  
EDI  
Pro  
Str  
Esc  
AC  
My  
Ho  
Ho  
flo  
as  
25  
Un