

1





Calcium Polysulfide, not less the

INERT INGREDIENTS:

Not more than.

For the Treatment of Psoroptic and Sarcoptic Scabies in Cattle, Horses, Sheep, and Swine. Also Effective for Sarcoptic Mange of Dogs and Cats.

B

DIRECTIONS

Dilute LIME & SULFUR SOLUTION with warm water in the following proportions and use as a dip or spray:

Horses and Cattle_____ Sheep, Dogs, and Cats_____ Swine_____

or when hand treated_

Maintain the solution between 95° and 105° F. Use the solution but once and re-treat animals at 12 day intervals. Treatment may be repeated at 3 to 7 day intervals on dogs and when hand treating swine. CAUTION

KEEP OUT OF REACH OF CHILDREN

Wash hands after use. Do not contaminate food or water. Harmful if swallowed.

U.S.D.A. REG. NO. 1112 - 4

Distributed By

L. A. MOSHER CO. ATLANTA, GA. .

	30%
	. 70%
•	100%

 		_ 1	to	15
 		_ 1	to	20
 1	to	15	or	20,
 <u>.</u>		_ 1	to	10





المنبع ال

ONE GALLON

LIME & SULFUR SOLUTION

ACTIVE INGREDIENT:

Calcium Polysulfide, not less than 30% Not more than. 70% 100% For the Treatment of Psoroptic and Sarcoptic Scabies in Cattle.

INERT INGREDIENTS:

Horses, Sheep, and Swine. Also Effective for Sarcoptic Mange of Dogs and Cats.

DIRECTIONS

Dilute LIME & SULFUR SOLUTION with warm water in the following proportions and use as a dip or spray: Horses and Cattle_____ 1 to 15 Swine_____1 to 15 or 20,

or when hand treated_____1 to 10

Maintain the solution between 95° and 105° F. Use the solution but once and re-treat animals at 12 day intervals. Treatment may be repeated at 3 to 7 day intervals on dogs and when hand treating swine. CAUTION

> KEEP OUT OF BEACH OF CHILDREN water. Humild a swillow st.


Walb Linds offer say Do not retaining for Lor

U.S.D.A. REG. NO. 1112 - 4

Distributed By

L. A. MOSHER CO. ATLANFA, GA. -





ONE GALLON

LIME & SULFUR SOLUTION

ame

ACTIVE INGREDIENT:

Calcium Polysulfide, not less than.....

INERT INGREDIENTS:

Not more than.

For the Treatment of Psoroptic and Sarcoptic Scabies in Cattle. Horses, Sheep, and Swine. Also Effective for Sarcoptic Mange of Dogs and Cats.

DIRECTIONS

Dilute LIME & SULFUR SOLUTION with warm water in the following proportions and use as a dip or spray:

owing j	propo.				 	-	-	
Horses	and	Cattle	3		 		_ 1	
Sheep,	Dogs,	and	Cats		 		_ 1	
Swine.		·······			 1 1	to	15	(
or	when	hand	treated	· · · · · · · · · · · · · · · · · · ·	 		_ 1	

Maintain the solution between 95° and 105° F. Use the solution but once and re-treat animals at 12 day intervals. Treatment may be repeated at 3 to 7 day intervals on dogs and when hand CAUTION treating swine.

KEEP OUT OF REACH OF CHILDREN

Wash hands after use. Do not containnate lood or water. Humful if swallowed.

U.S.D.A. REG. NO. 1112 - 4

Distributed By

L. A. MOSHER CO. ATLANTA, GA. .

····	70	%
-	_	

100%

to 15 to 20

or 20, to 10

