

WARNING

May be fatal if swallowed. Do not get on skin. In case of contact, wash immediately with soap and water. Do not breathe vapor or spray mist. May cause poisoning by inhalation or skin absorption. Do not dispose of any unused material in such a manner that contact with humans or beneficial animals may occur. Do not contaminate feed and foodstuffs. Use other than directed may contaminate the crop.

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

ANTIDOTE: *External*—In case of spillage on skin, wash immediately with soap and water. *Internal*—If swallowed, call a physician immediately. Unless instructed otherwise, induce vomiting by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of warm water).

DISPOSAL OF EMPTY CONTAINERS AND SURPLUS INSECTICIDE

Do not re-use containers. Take empty containers to a supervised sanitary land-fill dump, otherwise crush containers (if possible) and bury in an isolated place where they will not contaminate water supply. Bury any unused insecticide at least 18 inches deep in an isolated location where it will not contaminate water supplies.

COMBUSTIBLE. Do not use or store near heat or open flame.


KEEP FROM FREEZING

NON-WARRANTY NOTICE

Follow directions carefully. Seller warrants such directions to be proper and that product conforms to the descriptions on this label. Timing and method of application, weather and crop conditions, and other influencing factors in the use of the product are beyond the control of the Seller, so Seller makes no other warranty, express or implied (including a warranty of merchantability), with reference to the results or effects of use. No person has authority to make any representation not contained on this label.

U.S. A. REG. NO. 114574

AMOCO Heptachlor 3 EC Emulsifiable Concentrate



AMERICAN CYANAMID COMPANY

WARNING KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL

1 gallon of AMOCO Heptachlor 3 EC contains 3 pounds of Heptachlor

DIRECTIONS FOR USE

AMOCO Heptachlor 3 EC is always diluted before application by emulsifying with water. Use 15 to 35 gallons spray per acre in conventional ground spray equipment. Recommendations given are for ground application equipment.

Consult State Agricultural Extension Service for additional information consistent with label directions and cautions.

ACTIVE INGREDIENTS by weight:	
Heptachlor	32.40%
Related compounds	11.41%
Aromatic petroleum solvent	56.19%
INERT INGREDIENTS by weight	
Heptachlor 3 EC (3.00 lb. Heptachlor per gallon)	5.00%
TOTAL	100.00%

Control of Soil Insects on Corn

Soil Insect Complex—Born resistant Corn Rootworms, Seed Corn Beetle, Seed Corn Maggot, Wireworm, Grape Crickets, Cutworms. For broadcast application mix 4 to 5 1/2 pints AMOCO Heptachlor 3 EC (1 1/2 to 2 pounds Heptachlor) in enough water to cover one

acre. Use 1/2 pint per acre for broadcast application. For row application use 2 1/2 pints AMOCO Heptachlor 3 EC per acre. Heptachlor 3 EC is not for use on corn. Before any treatment, consult your local extension agent. Wireworms are not a pest on corn. Recommendations for 22 row and 4

Corn Crickets—Use 1/2 pint per acre for broadcast application. State authorities use 1/2 pint AMOCO Heptachlor 3 EC per acre in sufficient water to treat one acre row spraying. For broadcast treatment use 1 to 1 1/2 pints Heptachlor 3 EC per acre. Mix into a 2 to 4 inches of soil after broadcast application. In Missouri other states—check state recommendations. 1/2 pint per acre broadcast.

Cutworms—Use 2 1/2 to 5 pints (1 to 2 1/2 pints per acre) as a broadcast treatment preferably 10 to 14 days before planting. 1 to 4 inches immediately after application a minimum average rate of 4 pints is recommended.

Wireworms—Use 4 to 5 pints (2 1/2 to 4 pints per acre) as a broadcast treatment preferably 10 to 14 days before planting. Work into top 1 to 4 inches of soil. Use 3 pints per acre for high population of larvae.

White Grubs—Use 4 to 5 pints (2 1/2 to 4 pints per acre) as a broadcast treatment before planting and immediately after.

Seed Corn Beetle and Seed Corn Maggot—Use 5 1/2 pints (3 1/2 to 4 pints) Heptachlor 3 EC per acre as a broadcast application. Use 1/2 pint per acre as a broadcast application.

Grasshoppers

Grasshoppers on non-cropped land, Fence Rows, Ditch Banks, etc.—Use 1/2 pint AMOCO Heptachlor 3 EC per acre (1/2 pint per acre). Treat fence rows, ditch banks, cropped land. Do not graze dairy or beef treated areas. Do not feed forage from dairy or meat animals. Birds feeding on treated areas may be killed.

MADE IN U.S.A.

Net Contents


U.S. GALS.

AMOCO

Heptachlor

3 EC

Emulsifiable Concentrate



PRODUCT OF AMERICAN OIL COMPANY

WARNING KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL

1 gallon of AMOCO Heptachlor 3 EC
contains 3 pounds of Heptachlor

DIRECTIONS FOR USE

AMOCO Heptachlor 3 EC is always diluted before application by emulsifying with water. Use 15 to 35 gallons spray per acre in conventional ground spray equipment. Recommendations given are for ground application equipment.

Consult State Agricultural Extension Service for additional information consistent with label directions and cautions.

ACTIVE INGREDIENTS (by weight)	
Heptachlor	32.49%
Related compounds	11.41%
Aromatic petroleum solvent	51.10%
INERT INGREDIENTS (by weight)	
Heptachlor solvent	5.00%
TOTAL	100.00%

Control of Soil Insects on Corn

Soil Insect Complex Non-resistant Corn Rootworms, Seed Corn Beetle, Seed Corn Maggot, Wireworm, Grape Celaspis, Cutworms. For broadcast application mix 4 to 5½ pints AMOCO Heptachlor 3 EC (1½ to 2 pounds Heptachlor) in enough water to cover one

acre. Do not work into top 2 to 4 inches of soil immediately after application. For row crop treatment mix 2½ to 4 pints AMOCO Heptachlor 3 EC (¾ to 1½ pounds Heptachlor) in enough water to treat one acre. Broadcast treatment is preferred when Cutworms and Wireworms are the main problem. See specific recommendations for Wireworm and Cutworm control.

Corn Rootworm Non-resistant species and when such are reported by State authorities. Mix 4 to 5½ pints AMOCO Heptachlor 3 EC (1½ to 2 pounds Heptachlor) in sufficient water to treat one acre for row and row spraying. For broadcast treatment mix 2½ to 4 pints (¾ to 1½ pounds) Heptachlor with water to treat one acre. Mix into top 2 to 4 inches of soil immediately after broadcast application. In Missouri and possibly other states—check state recommendations—use 2 pints row, or 4 pints broadcast.

Cutworms—Use 2½ to 5½ pints (¾ to 2 pounds Heptachlor) per acre as a broadcast treatment applied to soil preferably 10 to 14 days before planting. Mix into top 3 to 4 inches immediately after application. In some states a minimum dosage rate of 4 pints (1½ pounds) is recommended.

Wireworms—Use 4 to 6 pints (1½ to 2 pounds) Heptachlor per acre as a pre-plant broadcast soil application preferably 10 to 14 days before planting. The granules work into top 3 to 4 inches of soil. Use 6 pint dosage on peat soils or for high population of large worms.

White Grubs—Use 5 to 8 pints (2 to 3 pounds) Heptachlor per acre as a broadcast treatment 10 to 14 days before planting and immediately disperse.

Seed Corn Beetle and Seed Corn Maggot—Use 4 to 5½ pints (1½ to 2 pounds) Heptachlor per acre as a pre-plant broadcast application or use 1½ to 2½ pints applied in row at planting time.

Grasshoppers

Grasshoppers on non-cropped land, such as Fence Rows, Ditch Banks, etc.—Use 1½ pints of AMOCO Heptachlor 3 EC per acre. 2 gallons Heptachlor. Treat fence rows, ditch banks, etc. other non-cropped land. Do not graze dairy or meat animals on treated areas. Do not feed crop from treated areas. Dairy or meat animals feeding on treated areas may be killed.

MADE IN U.S.A.

AMOCO 1961

Net Contents

U.S. GALS.