

DOSAGE and ADMINISTRATION: Shake well before using. Apply with a swab or soft bristle brush. Fresh wounds should be protected by a thorough coating of Myzin Smear. Infested wounds should be well packed with the smear and coating placed completely around the wound so it covers all blood and exudate. Repeat application twice a week until worm-infested wound is healed.

PRECAUTIONS: Not effective in some populations or strains of screwworms which are resistant to Lindane. This will be observed by lack of response to treatment.

E.P.A. Reg. No. 1117-24



381

Myzin® Smear

For control of screwworms in horses.

Active Ingredients:

Gamma isomer of benzene hexachloride from Lindane	3% w/w (3.4 Gm.)
Pine oil	35% w/w (39.7 Gm.)
Inert Ingredients	62% w/w
Total	100% w/w

CAUTION: Keep out of reach of children. See side panel for other cautions.

NET WT. 4 OZ.

Fort Dodge Laboratories, Inc.
Fort Dodge, Iowa 50501
Sales to Veterinarians Only

WARNING: Use sparingly on newborn (very young) individuals (never in amount to exceed 1 or 2 level teaspoonfuls at one time on newborns). Do not use on dairy animals or other animals intended for use for food. Do not use on dogs, cats or other household pets.

CAUTION: Harmful if swallowed. If swallowed, do not induce vomiting. Vomiting may be harmful. Call a physician immediately. Avoid breathing of vapor. May be absorbed through skin. Do not get in eyes or on skin. If contact occurs with eyes, flush immediately with water. In case of spillage on skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not re-use containers. Destroy when empty.

Made in U.S.A.
3863A

Lot

ACCEPTED

FORT DODGE

MYZIN® SMEAR

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND ROUDICIDE ACT FOR ECONOMIC POISON REGISTER-
E.P.A. REG. NO. 1117-24

Active ingredients:

Gamma isomer of benzene hexachloride from Lindane	3% w/w (108.86 Gm.)
Pine oil	35% w/w (1.27 Kg.)
Inert ingredients	62% w/w
Total	100% w/w

DOSE and ADMINISTRATION: SHAKE WELL BEFORE USING. Apply with a swab or soft bristle brush. Fresh wounds should be protected by a thorough coating of Myzin Smear. Infested wounds should be well packed with the smear and coating placed completely around the wound so it covers all blood and exudate. Repeat application twice a week until worm-infested wound is healed.

PRECAUTIONS: Not effective in some populations or strains of screwworms which are resistant to Lindane. This will be observed by lack of response to treatment.

WARNING: Use sparingly on newborn (very young) individuals (never in amount to exceed 1 or 2 level teaspoonfuls at one time on newborns). Do not use on dairy animals or other animals intended for use for food. Do not use on dogs, cats or other household pets.

CAUTION: Keep out of reach of children.

Harmful if swallowed. If swallowed, do not induce vomiting. Vomiting may be harmful. Call a physician immediately. Avoid breathing of vapor. May be absorbed through the skin. Do not get in eyes or on skin. If contact occurs with eyes, flush immediately with water. In case of spillage on skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not reuse container. Destroy when empty.

E.P.A. Reg. No. 1117-24

NET WT. 8 LB.

Lot

Fort Dodge Laboratories, Inc.

3864A
Made in U.S.A.

Fort Dodge, Iowa 50501
Sales to Veterinarians Only

DOSAGE and ADMINISTRATION: Shake well before using. Apply with a swab or soft bristle brush. Fresh wounds should be protected by a thorough coating of Myzin Smear. Infested wounds should be well packed with the smear and coating placed completely around the wound so it covers all blood and exudate. Repeat application twice a week until worm-infested wound is healed.

PRECAUTIONS: Not effective in some populations or strains of screwworms which are resistant to Lindane. This will be observed by lack of response to treatment.

E.P.A. Reg. No. 117-24



Myzin® Smear

For control of screwworms in horses.

Active Ingredients:

Gamma isomer of benzene hexachloride from Lindane	3% w/w (3.4 Gm.)
Pine oil	35% w/w (39.7 Gm.)
Inert Ingredients:	62% w/w
Total	100% w/w

CAUTION: Keep out of reach of children. See side panel for other cautions.

NET WT. 4 OZ.


Fort Dodge Laboratories, Inc.
Fort Dodge, Iowa 50501
Sales to Veterinarians Only

WARNING: Use sparingly on newborn (very young) individuals (never in amount to exceed 1 or 2 level teaspoonfuls at one time on newborns). Do not use on dairy animals or other animals intended for use for food. Do not use on dogs, cats or other household pets.

CAUTION: Harmful if swallowed. If swallowed, do not induce vomiting. Vomiting may be harmful. Call a physician immediately. Avoid breathing of vapor. May be absorbed through skin. Do not get in eyes or on skin. If contact occurs with eyes, flush immediately with water. In case of spillage on skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not re-use containers. Destroy when empty.

Made in U.S.A.
3863A

Lot



Active ingredients:

Gamma isomer of benzene hexachloride from Lindane	3% w/w (108.86 Gm.)
Pine oil	35% w/w (1.27 Kg.)
Inert ingredients:	62% w/w
Total	100% w/w

DOSE and ADMINISTRATION: SHAKE WELL BEFORE USING. Apply with a swab or soft bristle brush. Fresh wounds should be protected by a thorough coating of Myzin Smear. Infested wounds should be well packed with the smear and coating placed completely around the wound so it covers all blood and exudate. Repeat application twice a week until worm-infested wound is healed.

PRECAUTIONS: Not effective in some populations or strains of screwworms which are resistant to Lindane. This will be observed by lack of response to treatment.

WARNING: Use sparingly on newborn (very young) individuals (never in amount to exceed 1 or 2 level teaspoonfuls at one time on newborns). Do not use on dairy animals or other animals intended for use for food. Do not use on dogs, cats or other household pets.

CAUTION: Keep out of reach of children. Harmful if swallowed. If swallowed, do not induce vomiting. Vomiting may be harmful. Call a physician immediately. Avoid breathing of vapor. May be absorbed through the skin. Do not get in eyes or on skin. If contact occurs with eyes, flush immediately with water. In case of spillage on skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not reuse container. Destroy when empty.

E.P.A. Reg. No. 117-24

NET WT. 8 LB.

Lot

Fort Dodge Laboratories, Inc.
Fort Dodge, Iowa 50501
Sales to Veterinarians Only

3864A
Made in U.S.A.

FORT DODGE

JAN 9 1974

MYZIN[®] SMEAR

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR CONTROL OF SCREW-WORMS IN HORSES
FOR ECONOMIC POISON REGISTER-

Active ingredients:

Gamma isomer of benzene hexachloride from Lindane.	3% w/w (108.86 Gm.)
Pine oil	35% w/w (1.27 Kg.)
Inert ingredients:	62% w/w
Total	100% w/w

DOSE and ADMINISTRATION: SHAKE WELL BEFORE USING. Apply with a swab or soft bristle brush. Fresh wounds should be protected by a thorough coating of Myzin Smear. Infested wounds should be well packed with the smear and coating placed completely around the wound so it covers all blood and exudate. Repeat application twice a week until worm-infested wound is healed.

PRECAUTIONS: Not effective in some populations or strains of screwworms which are resistant to Lindane. This will be observed by lack of response to treatment.

WARNING: Use sparingly on newborn (very young) individuals (never in amount to exceed 1 or 2 level teaspoonfuls at one time on newborns). Do not use on dairy animals or other animals intended for use for food. Do not use on dogs, cats or other household pets.

CAUTION: Keep out of reach of children.

Harmful if swallowed. If swallowed, do not induce vomiting. Vomiting may be harmful. Call a physician immediately. Avoid breathing of vapor. May be absorbed through the skin. Do not get in eyes or on skin. If contact occurs with eyes, flush immediately with water. In case of spillage on skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not reuse container. Destroy when empty.

E.P.A. Reg. No. 1117-24

NET WT. 8 LB.

Lot

Fort Dodge Laboratories, Inc.

3864A
Made in U.S.A.

Fort Dodge, Iowa 50501
Sales to Veterinarians Only