

ACCEPTED

AUG 30 1956

INSECTICIDE
FEDERAL INSECTICIDE ACT
EPA REG. NO. 1117-30One
Gallon

FORT DODGE

NOLVASAN *

(Chlorhexidine)

SOLUTION

COMPOSITION

Active ingredient:

Hexamethylenebis (p-chloro-phenyl-diguanide)

diacetate..... 2%

Inert ingredients: 98%

General Directions for Disinfection

After thorough washing, rinse the containers or utensils in a solution made by adding 1 tablespoonful of Nolvasan to each gallon of clean water. The cap of this bottle may be used for measuring. 1 capful being sufficient for each gallon of water.

For aiding in the control of virus diseases, such as infectious bronchitis or canine distemper - clean the walls, cages or utensils thoroughly and spray the surfaces with a diluted Nolvasan solution by adding 1 or 2 tablespoonfuls, or capfuls, to each gallon of clean water.

Cloths used for udder cleansing should remain in the disinfectant solution for 2 minutes before applying. This will aid in the control of mastitis.

Caution: Nolvasan in its concentrated form may be irritating to eyes, skin or mucous membranes. Wash exposed tissues thoroughly. Avoid contamination of feed or foodstuffs.

FORT DODGE LABORATORIES, INC.

Fort Dodge, Iowa

Sales to Veterinarians Only

Made in U.S.A.

* Trade Mark