

NOLVASAN SOLUTION
(Chlorhexidine)

Dairy Sanitizing - General Directions


After thorough washing, rinse the teatcups, containers, other utensils or milking lines with a solution made by adding 1 tablespoonful to each gallon of clean water. Parts should remain in contact with the sanitizing solution for 2 minutes.

Cloths used for udder cleansing should also remain in the sanitizing solution for 2 minutes before re-use. This will aid in the control of mastitis. The ends of the teats may be dipped in the solution to prevent chapping and hasten healing of any wounds or abrasions.

Registered by U.S.D.A.
No. 1117-30

Made in U.S.A.

Sales to Veterinarians Only



ONE GALLON

NOLVASAN
(Chlorhexidine)
SOLUTION

Active Ingredients:

Hexamethylenebis (p-chloro-phenyl-diguanide)	
Diacetate	2%
Inert Ingredients	98%

General Directions for Disinfection

After thorough washing, rinse the containers or utensils in a solution made by adding 1 tablespoonful of Nolvasan to each gallon of clean water. The cap of this bottle may be used for measuring, 1 capful being sufficient for each gallon of water.

For aiding in the control of virus diseases, such as infectious bronchitis or canine distemper--clean the walls, cages or utensils thoroughly and spray the surfaces with a diluted Nolvasan solution by adding 1 or 2 tablespoonfuls, or capfuls, to each gallon of clean water.

CAUTION: NOLVASAN SOLUTION may be irritating to eyes or mucous membranes.

Manufactured by
FORT DODGE LABORATORIES, INC.
Fort Dodge, Iowa

by arrangement with
Imperial Chemical Industries, Ltd.
Pharmaceuticals Division

1117-30

03/15/1957

1/1