1 gallon



(CHLORHEXIDINE)

Virucide *

Active ingredient:
1, 1'-Hexamethylenebis [5-(p-chlorophenyl) biguanide] diacetate 2% . . . 98% Inert ingredients

General Directions for Disinfection

After thorough washing, rinse the containers or utensils in a solution made by adding 1 ounce (2 tablespoonfuls) of Nolvasan solution to each gallon of clean water. Not effective against gram-positive pyogenic cocci.

For disinfection of inanimate objects to aid in control of Newcastle virus, canine distemper virus, infectious bronchitis virus, bovine rhinotracheitis virus, bovine virus diarrhea virus and parainfluenza-3 virus—clean walls, cages or utensils thoroughly and spray or wash surface with a solution made by adding 3 ounces (6 tablespoonfuls) of Nolvasan Solution to each gallon of clean water.

CAUTION: Keep out of reach of children. May be irritating to eyes or mucous membranes. Avoid contamination of feed and foodstuffs.

Made in U.S.A. by arrangement with Imperial Chemical Industries, Limited

Fort Dodge Laboratories, Inc.

Fort Dodge, Iowa Sales to Veterinarians Only lated company U.S.D.A. Reg. No. 1117-30 HFG *8619,354 of related company

ACCEPTED 12/13/67