

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

##### Veterinary or Farm Premises

- 1 Remove all animals and feed from premises, vehicles, and other equipment.
- 2 Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.
- 3 Empty all troughs, racks, and other feeding and watering appliances.
- 4 Thoroughly clean all surfaces with soap or detergent and rinse with water.
- 5 Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
- 6 Invert all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
- 7 Ventilate buildings, vehicles, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried.
- 8 Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

##### For disinfection of official meat, poultry, rabbit and egg establishments

- 1 All food products and packaging material must be removed from the room or carefully covered and protected.
- 2 Remove any loose dirt, litter, etc. that might be lying on floor or attached to the equipment.
- 3 Thoroughly clean all surfaces with soap or detergent and rinse with water.
- 4 Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
- 5 Expose or soak all equipment and/or utensils with the recommended disinfecting solution for a period of 10 minutes.
- 6 After disinfection all equipment and/or utensils must be thoroughly rinsed with potable water before operations are resumed.

For dipping teats as an aid in controlling bacteria that causes mastitis  
Immediately after the cow is milked dip each teat into the dipping solution. Teat dipping should start one week before the cow breeds. When drying off a cow the teats should continue to be dipped once a day for 3 to 4 days. Under and teats of the cow must be thoroughly washed before milking.

#### FRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

##### CAUTION:

May be harmful if swallowed. May be irritating to eyes or mucous membranes. In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists.

##### Environmental Hazards:

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### STORAGE AND DISPOSAL

PROHIBITION: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Disinfectant mixture or waste that can not be used or chemically reprocessed should be disposed of in a landfill approved for disinfectants or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose in an incinerator or land fill approved for disinfectant containers or bury in a safe place.

GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

0626D

Fort Dodge Laboratories, Inc.

Fort Dodge, Iowa 50501

EPA Est. No. 1117-141, EPA Reg. No. 1117-30

ACCEPTED  
by CHAMBERS  
EPA Letter Dated

AUG 24 1963

Under the authority of the Federal Insecticide, Fungicide and Rodenticide Act, as amended, this product is registered for use in the United States.  
Registered under EPA Reg. No. 1117-30  
442-50

THIS DOCUMENT AVAILABLE

**BEST DOCUMENT AVAILABLE**

**DIRECTIONS FOR USE  
GENERAL CLASSIFICATION**

It is a violation of Federal Law to sell this product or its contents without labeling.

**Foaming Insurance** is a concentrated disinfectant for institutional use. It is an industrial cleaning agent used in Nursing Homes, Hospitals, Schools, Laboratories, Institutional Facilities and Factories.

**DISINFECTING DEODORIZING  
AND CLEANING**

To disinfect hard surfaces such as bathtubs, tile, walls, shelves, tables, chairs and desks. Shake can before use. Spray on surface to be treated. Let stand for 10 minutes. Wipe off with clean white cloth and rinse with water. Scrubbing is recommended when Foaming Insurance is used on heavily soiled areas.

**MILDEWSTAT** - To control mold and mildew growth. Apply to walls, tile, floors, fixtures, etc. Rinse with water.

**PRECAUTIONARY STATEMENT  
HAZARDS TO HUMAN  
AND DOMESTIC ANIMALS  
WARNING**

For Industrial and Institutional Use Only. Do not drink. Do not breathe spray mist. Avoid contact with eyes.

**STATEMENT OF  
PRACTICAL TREATMENT**

Do not drink. If swallowed, do not induce vomiting. Call physician. If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water.

**NOTE TO PHYSICIAN**

**FOAMING  
INSURANCE**

**CLEANER  
DEODORIZER  
AEROSOL  
DETERGENT  
DISINFECTANT**

For Industrial and Institutional Use Only.

CONTENTS: 12 oz. (355 ml.)  
NET WT. 12 oz. (355 ml.)  
MANUFACTURED BY:  
Vestal Chemical Company  
A Division of Vesta Corporation  
St. Louis, Missouri 63126

Only for Sale To Use and Storage  
by Service Personnel

**KEEP OUT OF REACH  
OF CHILDREN  
WARNING**

**Foaming Insurance** is formulated to neutralize up to a five quart fire volume. It is a fast acting antiseptic disinfectant. It kills 99.9% of bacteria. It kills fungi. Helps to control insects in Industrial Plants, Medical and Dental Offices, Veterinary Clinics, Motels, Restaurants and other public areas. When used as directed, **Foaming Insurance** is tested to clean and disinfect inanimate washable hard surfaces such as walls, sinks, tables, chairs, telephones, bath rooms, shower stalls, bathtubs, chandeliers, cabinets and floorings.

In addition, **Foaming Insurance** deodorizes those areas which are generally hard to keep free of smelling such as garbage cans, trash, garbage bins and cans, bathrooms, restrooms and other areas where people frequently gather.


Another important fact about **Foaming Insurance** is its extremely effective disinfecting and sterilizing qualities. This makes it ideal for medical and dental offices, hospitals, restaurants, laundries, food processing plants, etc. Effectiveness has been established that **Foaming Insurance** is as effective as steam sterilization in the presence of organic materials.

**Foaming Insurance** is also an excellent barrier against mold and mildew. Tremendous strides have been made when used as directed. Mildewstat is used in the bathroom, laundry room, kitchen, etc. It effectively controls mold and mildew on surfaces from 10 to 12 feet.

**Foaming Insurance** is strong and potent. When diluted, it is extremely safe and non-toxic to all surfaces.

**STORAGE AND DISPOSAL**

Keep out of reach of children.  
Store in a cool dry place.



ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 24 1983

Under the Federal Insecticide,  
Fungicide and Rodenticide Act  
as amended, the product  
registered under EPA Reg. No.

1117 35



# NOLVASAN® SOLUTION

## NOLVASAN SOLUTION

### Recommended Concentration For Use

- I For disinfection of inanimate objects to aid in control of canine distemper virus, equine influenza virus, transmissible gastroenteritis virus, hog cholera virus, parainfluenza-3 virus, bovine rhinotracheitis virus, bovine virus diarrhea virus, infectious bronchitis, Newcastle virus, Venezuelan encephalitis virus, equine rhinopneumonitis virus, feline rhinotracheitis virus and pseudorabies virus—3 ounces (16 fl. oz. equivalent) per gallon of clean water. Nolvasan Solution has been shown to be virucidal in vitro against rabies virus (CVS strain) in laboratory tests when used as directed above.
- II For control of veterinary external parasites—1/2 oz. (2 tablespoons—1/4 cup) per gallon of clean water.
- III For disinfection of clothing, poultry, rabbit housing establishments—1 ounce (2 tablespoons—1/4 cup) of Nolvasan Solution to each quart of clean water.
- IV For dipping tears as an aid in controlling bacteria that causes eye virus. Make up a 1% dip by diluting 100 ml. of Nolvasan Solution to 10 liters of clean water. Use 10 ml. of this dip per eye and then add 10 ml. of tap water until you have a total volume of 100 ml.

Not effective against Pseudomonas aeruginosa or gram positive bacteria on mucous surface.\*

\*According to A.O.A.C. Use Dilution Test Method

### Disinfectant

Active Ingredient		
1,1-Hexamethylenebis[5-(p-chlorophenyl)]	2%	
biquanide diacetate	98%	
Inert Ingredients		100%

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**

SEE SIDE PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS

1 GALLON

REST ROOM AVAILABLE