

1 gallon



MALINGO*

For control of ticks, lice, fleas, and sarcoptic and demodectic mange of dogs.

Active Ingredients:

- Technical Chlordane** 4% w/w
- Gamma isomer of benzene hexachloride from Lindane 1% w/w
- Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) 1% w/w
- Petroleum solvent 58.5% w/w
- Inert ingredients 35.5% w/w
- Total 100% w/w

**EQUIVALENT to 2.4% Octachloro-4, 7-methanotetrahydroindane and 1.6% related compounds.

CAUTION -- Keep out of the reach of children. See side panel for additional cautions.

Fort Dodge Laboratories, Inc.
Fort Dodge, Iowa 50501
Sales to Veterinarians Only

USAGE: Add 3 ounces (6 tablespoonfuls) Malingo to 1 gallon of water. Animal should first be bathed with soap and water, rinsed and toweled. Apply the diluted Malingo solution with sponge or cloth or by dipping, and allow the animal to dry without toweling. To prevent reinfestation repeat application every 2 to 3 weeks.

Since diagnosis of demodectic mange is often difficult, and treatment may involve other factors such as diet, treatment should be done by or under direction of a veterinarian.

CAUTION: FOR EXTERNAL USE ONLY. Use only the diluted solution as recommended. Harmful if swallowed. Avoid inhalation or contamination of feed. Repeated or prolonged contact with skin can cause toxic symptoms. Wash thoroughly after using. **DO NOT** use on cats, dairy cattle, nursing puppies, or animals being finished for slaughter.

Made in U.S.A.
DFI

*#678,281 of related company
U.S.D.A. Reg. No. 1117-31



AC...
JUL 18, 69

JUL 18 1969
UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND RODENT REPELLENT
FOR ECONOMIC PESTS IN REGISTRATION
ED UNDER NO. 1117-31 SUBJECT
TO ATTACHED COMMENTS.

WARNING

May be fatal if swallowed. Do not get on skin. In case of contact, wash immediately with soap and water. Do not breathe vapor or spray mist. May cause poisoning by inhalation or skin absorption. Do not dispose of any unused material in such a manner that contact with humans or beneficial animals may occur. Do not contaminate feed and foodstuffs. Use other than directed may contaminate the crop.

Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

ANTIDOTE: *External*—In case of spillage on skin, wash immediately with soap and water. *Internal*—If swallowed, call a physician immediately. Unless instructed otherwise, induce vomiting by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of warm water).

DISPOSAL OF EMPTY CONTAINERS AND SURPLUS INSECTICIDE

Do not re-use containers. Take empty containers to a supervised sanitary land-fill dump, otherwise crush containers (if possible) and bury in an isolated place where they will not contaminate water supply. Bury any unused insecticide at least 18 inches deep in an isolated location where it will not contaminate water supplies.

COMBUSTIBLE: Do not use or store near heat or open flame.

NON-WARRANTY NOTICE

Follow directions carefully. Seller warrants such directions to be proper and that product conforms to the descriptions on this label. Timing and method of application, weather and crop conditions, and other influencing factors in the use of the product are beyond the control of the Seller, so Seller makes no other warranty, express or implied (including a warranty of merchantability), with reference to the results or effects of use. No person has authority to make any representation not contained on this label.

AMOCO Heptachlor Spray

Emulsifiable Concentrate



PRODUCT OF AMERICAN OIL COMPANY
CHICAGO, ILL. USA 60680

WARNING KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL

ACTIVE INGREDIENTS (by weight)	
Heptachlor*	22.3%
Related compounds	8.6%
Aromatic petroleum solvent	63.1%
INERT INGREDIENTS (by weight)	
*Heptachlorotetrahydro-4,7-Methanoindene	6.0%
TOTAL	100.0%

DIRECTIONS FOR USE

AMOCO Heptachlor Spray is always diluted before application, by emulsifying with water.

Consult State Agricultural Extension Service for specific recommendations on timing and dosage rates in your area.

1 gallon of AMOCO Heptachlor Spray contains 2 pounds of Heptachlor

Control of Soil Insects on Corn

Soil Insect Complex (Non-resistant Corn Rootworms and Seed Corn Beetle, Seed Corn Maggot, Wireworm, Grape Colaspis, Cutworms)—For broadcast application mix 3 quarts AMOCO Heptachlor Spray (1½ pounds Heptachlor) in enough water to cover one acre. Do not work into top 3 to 4 inches of soil immediately after application. For band or row treatment mix 2 quarts Heptachlor Spray (1 pound Heptachlor) in sufficient

water to treat one acre. Broadcast treatment is where Curworms and/or Wireworms are present. See specific recommendations for Wireworm and Cutworm control.

Corn Rootworm (Non-resistant species and variety use is approved by State authorities)—Mix Heptachlor Spray (½ pound Heptachlor) in water to treat one acre for band or row spray. For broadcast treatment mix 2 quarts (1 pound Heptachlor) with water to treat one acre. Mix into top 3 to 4 inches of soil immediately after broadcast application.

Cutworms—Use 2 to 4 quarts (1 to 2 pounds Heptachlor) per acre as a broadcast treatment applied preferably 10 to 14 days before planting. Mix into top 3 to 4 inches immediately after application. State of Illinois states a minimum dosage rate of 3 quarts (1½ pounds Heptachlor) is recommended.

Wireworms—Use 3 to 5 quarts (1½ to 2½ pounds Heptachlor) per acre as a pre-plant broadcast application, preferably 10 to 14 days before planting. Thoroughly work into top 3 to 4 inches of soil. Use 4 quart dosage on peat soils or for high populations of large worms.

White Grubs—Use 4 to 6 quarts (2 to 3 pounds Heptachlor) per acre as a broadcast treatment 10 to 14 days before planting and immediately disced in.

Seed Corn Beetle (non-resistant) and Seed Corn Maggot—Use 3 to 4 quarts (1½ to 2 pounds Heptachlor) per acre as a pre-plant broadcast application. For row application use 1 to 2 quarts (½ to 1 pound Heptachlor) per row at planting time.

Grasshoppers

Grasshoppers on non-cropped land, such as Rows, Ditch Banks, etc.—Use 1 quart of Heptachlor Spray per acre. 2 quarts Heptachlor Spray per acre for ditch banks and other non-crop areas. Do not graze dairy or meat animals on treated areas. Do not feed forage from treated areas to dairy animals. Birds feeding on treated areas may

Net Co. Cents

U. S. GALS.