

WARNING: If swallowed by humans, induce vomiting by giving 2 tablespoons of salt in glass of water. Call physician immediately. Wash hands immediately after use. In case of accidental contact during administration, wash promptly and change clothes.

PHYSICIAN: Trichlorfon is a cholinesterase inhibitor. Atropine is an antidote. Give supportive treatment. Do not walk patient or give morphine.

CAUTION: Not to be used in horses intended for use as food. Store in cool, dry place.

0.76 oz.



Dyrex T.F. 200

Trichlorfon

+ Phenothiazine + Piperazine

Anthelmintic for Horses

CAUTION: Federal Law restricts this drug to sale by or on the order of a licensed veterinarian.

WARNING: Keep out of reach of children. See side panel for additional cautions.

FORT DODGE LABORATORIES, INC.
Fort Dodge, Iowa

For removal of large and small strongyles, bots, large roundworms (ascarids), and pinworms in horses.

This bottle contains:

Active Ingredients:
 0,0-dimethyl, 2,2,2-trichloro-1-hydroxyethyl phosphonate† 3.64 Gm. (16.8%)
 Phenothiazine, micronized .. 2.50 Gm. (11.5%)
 Piperazine dihydrochloride** 14.76 Gm. (68.2%)
 **Equivalent to 37% piperazine base or 8.0 Gm.
 Inert ingredients: 0.74 Gm. (3.5%)
 Total net contents:..... 21.64 Gm.

DOSAGE: For each 200 lb. bodyweight mix the contents of both this bottle and the inner vial in warm water and administer by stomach tube. Prepare each dose individually. Do not retain any unused suspension or powder for use at a later date. Do not fast horses before or after treatment. Symptoms of overdosage are ataxia and colic. Antidote: atropine. Mares heavy with foal should not be wormed.

It is important to read and understand product circular before using.

*Trademark of related company.
 †U. S. Patent 2,701,225, Canadian Patent 529,379, Chemagro Corp., licensee. Made in U.S.A.
 BFF U.S.D.A.Reg.No. 1117-37

ACCEPTED

APR 4 1968

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTER-
 ED UNDER NO. 14117-37 SUBJECT
 TO ATTACHED COMMENTS.

FORT DODGE

0.76 oz.

Dyrex T.F. 200

Trichlorfon
+ Phenothiazine + Piperazine

Anthelmintic for Horses

CAUTION: Federal Law restricts this drug to sale by or on the order of a licensed veterinarian.

WARNING: Keep out of reach of children. See side panel for additional cautions.

FORT DODGE LABORATORIES, INC.
Fort Dodge, Iowa

WARNING: If swallowed by humans, induce vomiting by giving 2 tablespoons of salt in glass of water. Call physician immediately. Wash hands immediately after use. In case of accidental contact during administration, wash promptly and change clothes.
PHYSICIAN: Trichlorfon is a cholinesterase inhibitor. Atropine is an antidote. Give supportive treatment. Do not walk patient or give morphine.
CAUTION: Not to be used in horses intended for use as food. Store in cool, dry place.

For removal of large and small strongyles, bots, large roundworms (ascarids), and pinworms in horses.

This bottle contains:

Active Ingredients:

0,0-dimethyl, 2,2,2-trichloro-1-hydroxyethyl phosphonate†	3.64 Gm. (16.8%)
Phenothiazine, micronized . . .	2.50 Gm. (11.5%)
Piperazine dihydrochloride**	14.76 Gm. (68.2%)
** Equivalent to 37% piperazine base or 8.0 Gm.	
Inert ingredients:	0.74 Gm. (3.5%)

Total net contents: 21.64 Gm.

DOSAGE: For each 200 lb. bodyweight mix the contents of both this bottle and the inner vial in warm water and administer by stomach tube. Prepare each dose individually. Do not retain any unused suspension or powder for use at a later date. Do not fast horses before or after treatment. Symptoms of overdosage are ataxia and colic. Antidote: atropine. Mares heavy with foal should not be wormed.

It is important to read and understand product circular before using.

†Trademark of related company.
‡U. S. Patent 2,701,225, Canadian Patent 529,379.
Chemagro Corp., licensee. Made in U.S.A.
BFF U.S.D.A.Reg.No. 1117-37

ACCEPTED

APR 4 1968

UNDER THE FEDERAL ANSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER. ED UNDER NO. 1117-37 SUBJECT TO ATTACHED COMMENTS.

WARNING: If swallowed by humans, induce vomiting by giving 2 tablespoons of salt in glass of water. Call physician immediately. Wash hands immediately after use. In case of contact during administration, wash promptly and change clothes.

PHYSICIAN: Trichlorfon is a cholinesterase inhibitor. Atropine is an antidote. Give supportive treatment. Do not walk patient or give morphine.

CAUTION: Not to be used in horses intended for use as food. Store in cool, dry place.

1.91 oz.



P305

Dyrex T.F. 500

Trichlorfon

+ Phenothiazine + Piperazine

Anthelmintic for Horses

CAUTION: Federal Law restricts this drug to use by or on the order of a licensed veterinarian.

WARNING: Keep out of reach of children. See side panel for additional cautions.

FORT DODGE LABORATORIES, INC.
Fort Dodge, Iowa 50501

For removal of large and small strongyles, bots, large roundworms (ascarids), and pinworms in horses.

This bottle contains:

Active Ingredients:

0,0-dimethyl, 2,2,2-trichloro-1-hydroxyethyl phosphonate†	9.10 Gm. (16.8%)
Phenothiazine, micronized ..	6.25 Gm. (11.5%)
Piperazine dihydrochloride**	36.90 Gm. (68.2%)
**Equivalent to 37% piperazine base or 20.0 Gm.	
Inert ingredients:	1.85 Gm. (3.5%)

Total net contents:..... 54.10 Gm.

DOSAGE: For each 500 lb. bodyweight mix the contents of both this bottle and the inner vial in warm water and administer by stomach tube. Prepare each dose individually. Do not retain any unused suspension or powder for use at a later date.

Do not fast horses before or after treatment. Symptoms of overdosage are ataxia and colic. Antidote: atropine.

Mares heavy with foal should not be wormed.

It is important to read and understand product circular before using.

*Trademark of related company.
†U. S. Patent 2,701,225, Canadian Patent 529,379,
Chemagro Corp., licensee. Made in U.S.A.
IFI U.S.D.A.Reg.No. 1117-38

Lot No.

Exp. Date

ACCEPTED WITH COMMENTS

HOW SUPPLIED

#200 bottle (one for each 200 lb. bodywt.).
#500 bottle (one for each 500 lb. bodywt.).
#1000 bottle (one for each 1,000 lb. bodywt.).

Packed in cartons of twelve (12).



DYREX* T. F.

(Trichlorfon + Phenothiazine + Piperazine)

Anthelmintic for Horses

This product has been specifically formulated as an equine anthelmintic and has been demonstrated to be active against all of the common equine parasites: ascarids, bots, large and small strongyles, and pinworms.

GENERAL INFORMATION

Preferably, all horses should be treated 1 month after the first killing frost and 1 month after being on pasture. Ascarids, pinworms and strongyles are usually removed within 3 days of treatment, bots by the 5th to 7th day.

A loosening of the stool may occasionally be seen several hours after treatment. Symptoms of overdosage are colic, diarrhea and ataxia, and appear within 1 to 3 hours after treatment. Such symptoms are usually transient and persist only briefly. Atrophine (5 to 10 mg./100 lb. bodyweight), administered to effect and repeated as necessary, is the antidote to aid in the control

Fort Dodge Laboratories, Inc.
Fort Dodge, Iowa 50501

*Trademark of related company.

Printed in U.S.A.

IFI

Jan., 1969

3 }
of these symptoms. Administer $\frac{1}{4}$ of dosage I.V. and remainder I.M. or S.C.

If desired, one of the oxime compounds which accelerate recovery of cholinesterase (such as 2-PAM, Protopam chloride) may be given in conjunction with the atropine.

PRECAUTIONS

Surgery or other severe stress should be avoided for at least 1 week before or after treatment. Increased side effects may be observed if the horse is being treated with a cholinesterase inhibitor, such as succinylcholine or organophosphorus or carbamate insecticides. Use of such agents should also be avoided for at least 1 week before and after treatment. Do not administer to sick, toxic or debilitated horses.

Field experience with Dyrex T.F. (trichlorfon-phenothiazine-piperazine) in pregnant animals would indicate no adverse effects. However, such treatment has historically been considered risky with any anthelmintic and, accordingly, treatment of mares in late pregnancy is not recommended.

ADMINISTRATION

The weight of each animal should be estimated carefully and the correct dosage prepared individually. The contents of both the outer bottle and the inner vial should be mixed together in warm water. Administration is by stomach tube.

To minimize the possibility of spillage, it is recommended that a funnel be used in pouring the suspension into the tube. After administration, the suspension remaining in the tube should be rinsed away with fresh water. Do not blow through the tube to empty. Flush unused suspension or powder down drain or bury.

DOSAGE

For removal of large and small strongyles, ascarids, bots and pinworms: Contents of one #200 bottle for each 200 lb. bodyweight; contents of one #500 bottle for each 500 lb. bodyweight; contents of one #1000 bottle for each 1,000 lb. bodyweight.

WARNING

Keep out of reach of children. If swallowed by humans, induce vomiting by giving 2 table-spoonfuls of salt in glass of water. Call physician immediately. Wash hands immediately after use. In case of contact during administration, wash promptly and change clothes.

PHYSICIAN: Trichlorfon is a cholinesterase inhibitor. Atropine is an antidote. Give supportive treatment. Do not walk patient or give morphine.

CAUTION

Federal law restricts this drug to use by or on the order of a licensed veterinarian. Not to be used in horses intended for use as food. Store in cool, dry place.