

WARNING: If swallowed by humans, induce vomiting by giving 2 tablespoons of salt in glass of water. Call physician immediately. Wash hands immediately after use. In case of accidental contact during administration, wash promptly and change clothes. **PHYSICIAN:** Trichlorfon is a cholinesterase inhibitor. Atropine is an antidote. Give supportive treatment. Do not walk patient or give morphine. **CAUTION:** Federal law restricts this drug to sale by or on the order of a licensed veterinarian. Not to be used in horses intended for use as food. Store in cool, dry place.



Dyrex T.F. 500

Trichlorfon
+ Phenothiazine + Piperazine
Anthelmintic for Horses

For removal of large and small Strongyles, Bots, large roundworms (Ascarids), and pinworms in horses.

FORT DODGE LABORATORIES, INC.
Fort Dodge, Iowa

Warning: Keep out of reach of children. See side panel for additional cautions.

This bottle contains:

Active Ingredients:

0,0-dimethyl, 2,2,2-trichloro-1-hydroxyethyl phosphonate†	9.10 Gm. (16.8%)
Phenothiazine, micronized . . .	6.25 Gm. (11.5%)
Piperazine Dihydrochloride**	36.90 Gm. (68.2%)
** Equivalent to 37% piperazine base or 20.0 Gm.	
Inert Ingredients	1.85 Gm. (3.5%)
Total net contents	54.10 Gm.

DOSAGE: For each 500 lb. bodyweight mix the contents of both this bottle and the inner vial in warm water and administer by stomach tube. Prepare each dose individually. Do not retain any unused suspension or powder for use at a later date. Do not fast horses before or after treatment. Symptoms of overdosage are ataxia and colic. Antidote: atropine. Mares heavy with foal should not be wormed.

It is important to read and understand product circular before using.

† Trademark of a Related Company.
‡ U. S. Patent 2,701,225, Canadian Patent 529,379,
Chemagro Corp., licensee. Made in U.S.A.
U.S.D.A.Reg.No. 1117-38

L 4614

ACCEPTED

NOV 5 1966

UNDER THE FEDERAL FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRATION UNDER NO. 1117-38