

A
No
UNDE
FUNGI
FOR E
ED UN
TO AT

WARNING: If swallowed by humans, induce vomiting by giving 2 tablespoons of salt in glass of water. Call physician immediately. Wash hands immediately after use. In case of accidental contact during administration, wash promptly and change clothes. **PHYSICIAN:** Trichlorfon is a cholinesterase inhibitor. Atropine is an antidote. Give supportive treatment. Do not walk patient or give morphine. **CAUTION:** Federal Law restricts this drug to sale by or on the order of a licensed veterinarian. Not to be used in horses intended for use as food. Store in cool, dry place.



Dyrex T.F. 1000

Trichlorfon
+ Phenothiazine + Piperazine

Anthelmintic for Horses

For removal of large and small Strongyles, Bots, large roundworms (Ascarids), and Pinworms in horses.

FORT DODGE LABORATORIES, INC.

Fort Dodge, Iowa

Warning: Keep out of reach of children. See side panel for additional cautions.

This bottle contains:

Active Ingredients:

0,0-dimethyl, 2,2,2-trichloro-1-hydroxyethyl phosphonate† 18.20 Gm. (16.8%)
Phenothiazine, micronized.. 12.50 Gm. (11.5%)
Piperazine Dihydrochloride** 73.80 Gm. (68.2%)

**Equivalent to 37% piperazine base or 40.0 Gm.

Inert Ingredients 3.70 Gm. (3.5%)

Total net contents.....108.20 Gm.

DOSAGE: For each 1000 lb. bodyweight mix the contents of both this bottle and the inner vial in warm water and administer by stomach tube. Prepare each dose individually. Do not retain any unused suspension or powder for use at a later date.

Do not fast horses before or after treatment. Symptoms of overdosage are ataxia and colic. Antidote: atropine.

Mares heavy with foal should not be wormed.

It is important to read and understand product circular before using.

†Trademark of a Related Company.

‡U. S. Patent 2,701,225, Canadian Patent 529,379, Chemagro Corp., licensee. Made in U.S.A.

U.S.D.A.Reg.No. 1117-39



EX T.F. 1000

Trichlorfon
thiazine + Piperazine
Anthelmintic for Horses

of large and small Strongyles,
roundworms (Ascarids), and
horses.

DODGE LABORATORIES, INC.
Fort Dodge, Iowa

Keep out of reach of children.
panel for additional cautions.

This bottle contains:

Active Ingredients:

0,0-dimethyl, 2,2,2-trichloro-1-hydroxyethyl phosphonate†	18.20 Gm. (16.8%)
Phenothiazine, micronized..	12.50 Gm. (11.5%)
Piperazine Dihydrochloride**	73.80 Gm. (68.2%)
**Equivalent to 37% piperazine base or 40.0 Gm.	
Inert Ingredients	<u>3.70 Gm. (3.5%)</u>
Total net contents.....	108.20 Gm.

DOSAGE: For each 1000 lb. bodyweight mix the contents of both this bottle and the inner vial in warm water and administer by stomach tube. Prepare each dose individually. Do not retain any unused suspension or powder for use at a later date. Do not fast horses before or after treatment. Symptoms of overdosage are ataxia and colic. Antidote: atropine. Mares heavy with foal should not be wormed.

It is important to read and understand product circular before using.

*Trademark of a Related Company.
†U. S. Patent 2,701,225, Canadian Patent 529,379,
Chemagro Corp., licensee. Made in U.S.A.
U.S.D.A.Reg.No. 1117-39

1117-39

ACCEPTED
NOV 5 1966
UNDER THE FEDERAL INSPECTION AND CONTROL ACT OF 1927
FOR FOOD AND DRUGS
ED UNDER THE
TO AT THE COMMENTS

FORT DODGE

Dyrex T.F.

Trichlorfon
+ Phenothiazine

Anthelmintic

For removal of large and
Bots, large roundworm,
Pinworms in horses.

FORT DODGE LABOR
Fort Dodge,

Warning: Keep out of r
See side panel for add

WARNING: If swallowed by humans, induce vomiting by giving 2 tablespoonfuls of salt in glass of water. Call physician immediately.

Wash hands immediately after use. In case of accidental contact during administration, wash promptly and change clothes.

PHYSICIAN: Trichlorfon is a cholinesterase inhibitor. Atropine is an antidote. Give supportive treatment. Do not walk patient or give morphine.

CAUTION: Federal Law restricts this drug to sale by or on the order of a licensed veterinarian. Not to be used in horses intended for use as food. Store in cool, dry place.