

CAUTION: Federal law restricts this drug to sale by or on the order of a licensed veterinarian. Not to be used in horses intended for use as food. Discard unused contents.

WARNING: If swallowed by humans, induce vomiting by giving 2 table spoonfuls of salt in glass of water. Call physician immediately. Wash hands immediately after use.

PHYSICIAN: This material is a cholinesterase inhibitor. Atropine is an antidote. Give supportive treatment. Do not walk patient or give morphine.

48.6
Gm


Dyrex
(trichlorfon)
Granules
Anthelmintic
for Horses
1000

Warning: Keep out of reach of children. See side panel for additional cautions.

This bottle contains:

Active ingredient:
0,0-dimethyl 2,2,2-trichloro-1-hydroxyethyl phosphonate 36.4 Gm. 75%
Inert ingredients 12.2 Gm. 25%

DOSAGE: For removal of Bots, Ascarids and Pinworms, give contents of one #1000 bottle for each 1,000 lb. bodyweight. For removal of Bots only, give one-half the regular dose.

Do not fast horses before or after treatment. Mix recommended dosage with a normal amount of grain. Remove and destroy any medicated feed not eaten within 12 hours. Mares heavy with foal should not be wormed.

Symptoms of overdose are ataxia and colic. Antidote: atropine.

It is important to read and understand product circular before using.

Fort Dodge Laboratories, Inc.
Fort Dodge, Iowa

Trademark of a Related Company.
U.S. Patent 2,701,725; Canadian Patent 529,379.
Chemagro Corp., licensee.

U.S.D.A. Reg. No. 1117-46

ACCEPTED
NOV 1961
61117-46