1 gallon



NOLVASAN-S

(CHLORHEXIDINE)

Virucide

Active ingredient: 1, 1'-Hexamethylenebis [5-(p-chlorophenyl) biguanide] diacetate 2% Inert ingredients98%

General Directions for Disinfection

After thorough washing, rinse the containers or utensils in a solution made by adding 1 outer 2 tablespoonfuls) of Nolvasan-S to each galon of clear water. Not effective against gram-positive pyogenic cocci. For disinfection of inalimate objects to aid in control of Newcastle virus, canine listempte virus, infectious bronchitis virus, bovine rhinor cheith virus, bovine virus diarrhea virus and parainfluenza-3 virus—clean walls, cages or utensils floroughly and pray or wash surface with a solution made by adding 3 ounces (6 tablespoonfuls) of Nolvasan-S to each gallon of clean water.

CAUTION: Keep out of reach of children. May be irritating to eyes or mucous membranes. Avoid contamination of feed and foodstuffs.

Made in U.S.A. by arrangement with
Imperial Chemical Industries, Limited

Fort Dodge Laboratories, Inc.

Fort Dodge, lowa L5769
Sales to Veterinarians Only
*®619,354 of related company HFG U.S.D.A. Reg. No. 1117-48

ACCEPTED

SEP1 4 1967

UNDER THE FEDERAL INSECTICIDE FUND HOUSE AND RODENTICIDE ACT FOR THE MAN POISON REGISTER-