

1117-48

03/22/1972

1/2

1 gallon

FORT DODGE®

M041

NOLVASAN-S®

(CHLORHEXIDINE)

Virucide†

Active Ingredients

1, 1'-Hexamethylenebis [3-(p-chlorophenyl)

biguanide] diacetate 2%

Inert Ingredients 98%

Directions

†For disinfection of inanimate objects to aid in the control of canine distemper virus, equine influenza virus, transmissible gastro-enteritis virus, hog cholera virus, parainfluenza-3 virus, bovine rhinotracheitis virus, bovine virus diarrhea virus, infectious bronchitis and Newcastle virus—after thorough cleaning, wash or rinse walls, cages or utensils with a solution made by adding 3 ounces (2 tablespoonfuls) of Nolvasan-S to each gallon of clean water.

Nolvasan-S has been shown to be virucidal in vitro against rabies virus (CVS strain) in laboratory tests when used as directed above.

For disinfection—After thorough cleaning, wash or rinse walls, cages or utensils with a solution made by adding 1 ounce (2 tablespoonfuls) of Nolvasan-S to each gallon of clean water. Nolvasan-S is not effective against *Pseudomonas aeruginosa* or gram-positive pyogenic cocci, such as *Staphylococcus aureus*.

CAUTION: Keep out of reach of children. May be irritating to eyes or mucous membranes. Avoid contamination of feed and foodstuffs.

Made in U.S.A. by arrangement with
Imperial Chemical Industries, Limited

Fort Dodge Laboratories, Inc.

Fort Dodge, Iowa 50501

Sales to Veterinarians Only
E.P.A. Reg. No. 1117-48

AGB

ACCEPTED

MAR 22 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT CIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 1117-48

Nolvasan "S"

One Pint



NOLVASAN "S"

(Chlorhexidine)

VIRUCIDE†

Active ingredient:

1,1'-Hexamethylenebis [5-(p-chlorophenyl)
biguanide] diacetate 2%

Inert ingredients: 98%

CAUTION: Keep out of reach of children. See back panel for other cautions.


Made in U.S.A. by arrangement with
Imperial Chemical Industries Limited

FORT DODGE LABORATORIES, INC.,

Fort Dodge, Iowa 50501

Sales to Veterinarians Only

Nolvasan "S"



DIRECTIONS

†For disinfection of inanimate objects to aid in the control of canine distemper virus, equine influenza virus, transmissible gastro-enteritis virus, hog cholera virus, parainfluenza-3 virus, bovine rhinotracheitis virus, bovine virus diarrhea virus, infectious bronchitis and Newcastle virus—after thorough cleaning wash or rinse walls, cages or utensils with a solution made by adding 3 ounces (6 tablespoonfuls) of Nolvasan-S to each gallon of clean water.

Nolvasan-S has been shown to be virucidal in vitro against rabies virus (CVS strain) in laboratory tests when used as directed above.

For disinfection — after thorough cleaning, wash or rinse walls, cages or utensils with a solution made by adding 1 ounce (2 tablespoonfuls) of Nolvasan-S to each gallon of clean water. Nolvasan-S is not effective against *Pseudomonas aeruginosa* or gram-positive pyogenic cocci, such as *Staphylococcus aureus*.

CAUTION: May be irritating to eyes or mucous membranes. Avoid contamination of feed or foodstuffs.

AGB

E.P.A. Reg. No. 1117-48