

PM 3, 1117-48

**ACCEPT**  
**NOV 03 19**  
 Under the Federal Insec  
 Fungicide, and Rodentic  
 as amended, for the pest  
 registered under  
 EPA Reg. No. 1117-

# **Nolvasan S**

Chlorhexidine diacetate

For cleaning, disinfection, deodorizing

## **Recommended Concentration For Use**

For disinfection of inanimate objects to aid in control of canine distemper virus, equine influenza virus, transmissible gastroenteritis virus, hog cholera virus, parainfluenza-3 virus, bovine rhinotracheitis virus, bovine virus diarrhea virus, infectious bronchitis virus, Newcastle virus, Venezuelan equine encephalitis virus, equine rhinopneumonitis virus, feline rhinotracheitis virus, pseudorabies virus, equine arteritis virus and canine coronavirus — 3 ounces (6 tablespoonfuls) per gallon of clean water. Nolvasan S has been shown to be virucidal *in vitro* against rabies virus (CVS strain) in laboratory tests when used as directed above.

For disinfection of veterinary or farm premises — 1 ounce (2 tablespoonfuls) per gallon of clean water.

Not effective against *Pseudomonas aeruginosa* or gram-positive cocci on inanimate surface.\*

\*According to A O A C. Use Dilution Test Method.

# **Nolvasan® S** Chlorhexidine diacetate

**Scented  
 Disinfectant**

**Bactericide Virucide†**



## **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

### **Veterinary or Farm Premises**

- 1 Remove all animals and feed from premises vehicles and other equipment.
- 2 Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals.
- 3 Empty all troughs, racks and other feeding and watering appliances.
- 4 Thoroughly clean all surfaces with soap or detergent and rinse with water.
- 5 Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
- 6 Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for moving litter and manure.
- 7 Ventilate buildings, vehicles and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried.
- 8 Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

## **Active Ingredient:**

1,1'-Hexamethylenebis [5-(p-chlorophenyl) biguanide] diacetate

## **Inert Ingredients**

2%  
 98%  
 100%

**KEEP OUT OF REACH OF CHILDREN**

## **CAUTION**

SEE SIDE PANEL FOR ADDITIONAL  
 PRECAUTIONARY STATEMENTS

**1 Pint**

**PRECAUTIONARY STATEMENTS  
 HAZARDS TO HUMANS  
 (AND DOMESTIC ANIMALS)**

## **CAUTION**

May be harmful if swallowed. May be irritating to eyes or mucous membranes. In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists.

### **Environmental Hazards**

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

## **STORAGE AND DISPOSAL**

**PROHIBITION:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Protect from freezing.

**PESTICIDE DISPOSAL:** Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Do not reuse containers. Rinse thoroughly with water and discard in trash.

**GENERAL:** Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

**Fort Dodge Laboratories, Inc.  
 Fort Dodge, Iowa 50501 USA**

EPA Est. No. 1117-IA-2 EPA Reg. No. 1117-48  
 91403 0405L

Lot  
 Exp. Date



**BEST AVAILABLE COPY**