

ACQUITTED
BY THE COURT
ON 12/10/57

1117-57

NOLVASAN® 6X
(Chlorhexidine acetate)

Recommended Concentration For Use


- I.* For disinfection of inanimate objects to aid in control of canine distemper virus, equine influenza virus, porcine transmissible gastroenteritis virus, hog cholera virus, bovine parainfluenza-3 virus, bovine rhinotracheitis virus, bovine virus diarrhea virus, avian infectious bronchitis, Newcastle virus, Venezuelan encephalitis virus, equine rhinopneumonitis virus, feline rhinotracheitis virus and pseudotuberculosis virus — ½ ounce (1 tablespoonful) per gallon of clean water.
- II. For disinfection of veterinary or farm premises — 5 ml. (1 teaspoonful) per gallon of clean water.
- III. For disinfection of official meat, poultry, rabbit and egg establishments — 5 ml. (1 teaspoonful) per gallon of clean water.

Not effective against *Pseudomonas aeruginosa* or gram-positive cocci on inanimate surface.**

* Pathogenic Fungi (*Trichophyton interdigitale*).

** According to A.O.A.C. Use Dilution Test Method.

M083



NOLVASAN®
6X
(Chlorhexidine acetate)

Bactericide
Virucide* Fungicide+

Active ingredient:	
1.1% — Hexamethylenebis [5-(p chlorophenyl) biguanide] acetate	12%
Inert ingredients	88%
	100%

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

1 GALLON

Panel 1 & 2 (One Gallon)

1117-57

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Veterinary or Farm Premises

1. Remove all animals and feed from premises, vehicles, and other equipment.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings, vehicles, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

For disinfection of official meat, poultry, rabbit and egg establishments

1. All food products and packaging material must be removed from the room or carefully covered and protected.
2. Remove any loose dirt, litter, etc., that might be lying on floor or attached to the equipment.
3. Thoroughly clean all surfaces with soap or detergent and rinse with water.
4. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
5. Expose or soak all equipment and/or utensils with the recommended disinfecting solution for a period of 10 minutes.
6. After disinfection all equipments and/or utensils must be thoroughly rinsed with potable water before operations are resumed.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION:

May be harmful if swallowed. May be irritating to eyes or mucous membranes. In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists.

Environmental Hazards

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

PROHIBITION: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Disinfectant, mixture or residue that can not be used or chemically reprocessed should be disposed of in a landfill approved for disinfectants or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for disinfectant containers, or bury in a safe place.

GENERAL: Consult Federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

0815A

Fort Dodge Laboratories, Inc.
Fort Dodge, Iowa 50501

EPA Est. No. 1117-IA-1 EPA Reg. No. 1117-57

Panel 3 & 4 (One Gallon)