DIRECTIONS

1. For disinfecting and cleaning in meat and poultry productions, such as in processing of meat, poultry, turkeys, and equipment, dilute Hockwald 1111 1 ounce to 1 gallon of vater (1:128). Wipe, mop or swab all surfaces with solution, leave in contact 10 minutes. All food contact surfaces must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.

2. For non-food disinfection of laying and broiler houses and AL INSECTICIDE hatcheries: Do not use in milking stalls, milking parlors or DENTICIDE ACT ilk houses. Remove all animals, poultry and feeds from the ISON REGISTER Premises. Remove all litter and manure from floors and other forms of the surfaces. Empty all feeding and watering troughs and applian-

buildings and other closed spaces. Do not house poultry, live-stock or employ equipment until treatment has been absorbed or dried. All treated feed racks, waterers and fountains must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.

3. For general cleanup in food processing plants, patient rooms, surgery, nursing homes, all animal facilities, on walls, floors, equipment, use 1 ounce Hockwald 1111 per 1 gallon of water (1:128). Cover surfaces thoroughly with solution and allow to remain in contact 10 minutes. For porous surfaces, increase concentration to 2 ounces per gallon. Rinse all surfaces coming in contact with food with potable water before reuse.

4. For disinfecting and deodorizing garbage and trash receptacles, carts and sick room equipment, use 1 ounce Hockwald 1111 to 1 gallon of water (1 part to 128 of water).

5. Hockwald 1111 is a prophylactic agent in controlling transmission of athlete's foot fungi on environmental surfaces. Prepare a solution of 1 ounce Hockwald 1111 per 1 gallon of water; mop or swab this solution onto floors and other surfaces normally contacted by the feet in locker rooms, shower stalls, bathrooms and around swimming pools.

AOAC USE DILUTION METHOD, 11TH EDITION

Staphylococcus aureus	1:128
Salmonella choleraesuis	1:128
Pseudomonasaeruginosa (PRD-10)	1:128

AOAC OFFICIAL FUNGICIDAL TEST, 11TH EDITION Trichophyton interdigitale 1:128

*Hockwald 1111 is a viricidal agent against Herpes simplex (strain HF) and influenza A₂/Aichi/2/68 (Hong Kong) viruses on inanimate environmental surfaces at 1:128 dilution.

Hockwald 1111 is Salmonellacidal - effective against the organisms which cause pullorum disease, fowl typhoid, and paratyphoid.

AOAC USE DILUTION METHOD, 11TH EDITION

Salmonella pullorum	1:128
Salmonella typhimurium	1:128
Salmonella typhosa	1.128
Salmonella choleraesuis	1:128

Hockwald 1111 is also effective against the following organisms using the AOAC USE DILUTION METHOD, 11TH EDITION.

1:128
1:128
1:128

AOAC TUBERCULOCIDAL TEST Mycobacterium tuberculosis

Hockwald 1111 removes mildew stains from hard surfaces.

1:128

Hockwald 1111 is conductive - may be used on conductive floorings as it does not interfere with electrical conductivity, as specified in the procedures in accordance with Paragraph 252, "Conductive Flooring of the National Board of Fire Underwriters," Specification No. 56.

DANGER

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Causes eye damage, and skin irritation. Do not get in eyes, on skin, or on clothing. Wash afterusing. Avoid contamination of food. Avoid inhalation of fogging mist. Do not use, pour, spill or store near heat or open flame.

FIRST AID: In case of contact, flush with water for 15 minutes and get medical attention, if irritation persists. For eyes flush thoroughly and get medical attention. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Call a physician immediately.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

Manufactured by

A Division of

OXFORD CHEMICAL

P.O. Box 227, Brisbane, Calif. 94005
7201 Paramount Blvd., Pico Rivera 90660

1974