

Not Reviewed. Registrations to be in accordance with
Federal Food, Drug, and Cosmetic Act, for Phase 2
of the Disinfectant, Antiseptic, and Sanitizer Act.

PUREX INSTITUTIONAL

FOR PROFESSIONAL SANITARY MAINTENANCE USE ONLY

NOT FOR SALE FOR RESIDENTIAL USE

Brillo PINOSAN

CLEANER • SANITIZER • DISINFECTANT - HOSPITAL TYPE

ACTIVE INGREDIENTS:

Sodium hypochlorite (5.25% available chlorine) 1.00
Sodium hydroxide (50% solution) 1.00
Sodium metasilicate (28% solution) 1.00
Sodium tetraborate (20% solution) 1.00
Sodium perborate (15% solution) 1.00
Sodium carbonate (50% solution) 1.00
Sodium silicate (28% solution) 1.00
Sodium borate (20% solution) 1.00
Sodium peroxide (15% solution) 1.00
Sodium hypochlorite (5.25% available chlorine) 1.00

INERT INGREDIENTS:

Water 1.00
Sodium hydroxide (50% solution) 1.00
Sodium metasilicate (28% solution) 1.00
Sodium tetraborate (20% solution) 1.00
Sodium perborate (15% solution) 1.00
Sodium carbonate (50% solution) 1.00
Sodium silicate (28% solution) 1.00
Sodium borate (20% solution) 1.00
Sodium peroxide (15% solution) 1.00
Sodium hypochlorite (5.25% available chlorine) 1.00

WARNING:

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

1 GALLON (3.78 LITERS)

ACTIVE INGREDIENTS:

N-alkyl 3,14,5,6; 3,12,4,5; 3,16,1,5,
dimethyl benzyl ammonium chloride 1.00

INERT INGREDIENTS: *

98.50

*This includes: sodium metasilicate and
sodium tetraborate, tetraacetic acid, tetraacetic salt

BEST AVAILABLE COPY

DIRECTIONS:

FOR CLEANING ONLY: Use Brillo Pinosan for all light to heavy duty cleaning on walls, woodwork, floors, in schools, court houses, hospitals, nursing homes, veterinarian clinics, industry, hotels, motels.

NORMAL SOIL: Mix 1 part Brillo Pinosan with 40-50 parts water.

LIGHT SOIL: Mix 1 part Brillo Pinosan with 60 parts water.

HEAVY SOIL: Mix 1 part Brillo Pinosan with 20 parts water.

TO CLEAN AND DISINFECT: Mix 1 part Brillo Pinosan with 20 parts water. Where *Pseudomonas aeruginosa* is suspected of being present, use 1 part Brillo Pinosan to 5 parts water.

WARNING: Avoid contamination of food stuffs. Harmful if swallowed. Not to be taken internally. May cause skin irritation or damage to the eyes. Do not get on skin or in eyes. In case of skin contact, flush with water. If irritation persists, obtain medical attention. For eyes, flush thoroughly with water and get medical attention.

A.O.A.C. USE DILUTION CONFIRMATION TEST

Salmonella choleraesuis (60 tubes) 1 to 20 *Escherichia coli* (60 tubes) 1 to 20
Staphylococcus aureus (60 tubes) 1 to 20 *Pseudomonas aeruginosa* (30 tubes) 1 to 5

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.



PUREX INDUSTRIAL
Turco Purex Industrial Corp.
7300 Bolsa Ave.
Westminster, CA 92684-3600

EPA Reg. No. 1124-57-AA

EPA Est. 1124-0H-3

45-7

F. R. PROFESSIONAL SANITARY

CLEA

ACTIVE INGREDIENTS
Sodium Hypochlorite 5.25%
Sodium Hydroxide 0.5%
Ammonium Hydroxide 0.5%
Formalin 0.5%
Sodium Chloride 0.5%
INEFFECTIVE INGREDIENTS

BEST AVAILABLE COPY

PA 2, 2