

Cross-infection is of major housekeeping concern not only in hospitals, but in schools, institutions, and industry. SANI-TURGE Disinfectant Cleaner is formulated for this problem area. It both cleans and disinfects effectively when used as directed. Its hard surface disinfecting action will reduce the hazard of cross-infection.

ACCEPTED
 2/24/72
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTER-
 ED UNDER NO. 1124-103

SANI-TURGE

DISINFECTANT CLEANER

Cleaner-Disinfectant-Deodorizer-Fungicide
 for Hospital and Institutional Use

WARNING: Keep out of reach of children. May cause severe eye irritation or eye damage. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

ANTIDOTES: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, give milk, raw egg white, gelatin solution; or if these are not available, drink water. Call a physician immediately.

ACTIVE INGREDIENTS:

Didecyl dimethyl ammonium chloride	4.25%
Tetrasodium ethylenediamine tetraacetate	1.60%
Sodium carbonate	2.00%
Sodium metasilicate, anhydrous	0.50%
INERT INGREDIENTS:	91.65%

WARNING: KEEP OUT OF REACH OF CHILDREN.
 See Left Panel for Additional Precautionary Statements.

55 GALLONS

Product Code F249C38

PUREX CORPORATION, I.T.D./INDUSTRIAL DIVISION
 24600 South Main Street
 Wilmington, California 90744

ACCEPTED

2/24/72

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
EPA UNDER NO. 1124-63

USE DIRECTIONS

Add two ounces per gallon water.

Apply SANI-TURGE Disinfectant Cleaner to walls, floors, and other hard surfaces such as tables, chairs and bed frames with a cloth or mop. For heavily soiled areas, a preliminary cleaning may be required.

At two ounces per gallon use-level, SANI-TURGE Disinfectant Cleaner is effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella choleraesuis*, and *Trichophyton interdigitale*. Germicidal performance against the first three organisms has been confirmed by the AOAC Use-Dilution Test. Fungicidal performance against *T. interdigitale* was determined by the AOAC Fungicidal Test.

Rinse empty container with water before discarding it.

AOAC PHENOL COEFFICIENTS

<i>Staphylococcus aureus</i> (ATCC 6538)	100.0
<i>Salmonella typhosa</i> (ATCC 6539)	43.8