

-2-

Note: The referral should be placed directly below the signal word.

f. Revise the statement:

...simplex Types I and II, Vaccinia...

to read:

...simplex Type 1 and 2, Vaccinia...

g. Add the following statement:


Only for Sale to, Use, and Storage by Service Persons.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

John M. Lee 
Product Manager (31)
Disinfectants Branch
Registration Division (TS-767C)

Enclosure

RD/DIS:JOB-91272:Lee:RD:85/02:Kendrick:898-1270:4/11/85:Del.4/18/85:EK

BEST AVAILABLE COPY

Precautionary Statements

Hazards to Humans and domestic animals

DANGER

Keep out of reach of children. Corrosive. Causes eye and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not breathe spray mist. Avoid contamination of feed.

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

STORAGE AND DISPOSAL

- DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

PESTICIDE DISPOSAL


PESTICIDE SPRAY MIXTURE OR RINSE WATER THAT CANNOT BE USED ACCORDING TO LABEL INSTRUCTIONS MUST BE DISPOSED OF ACCORDING TO FEDERAL OR APPROVED STATE PROCEDURES UNDER SUBTITLE C OF THE RESOURCE CONSERVATION AND RECOVERY ACT

CONTAINER DISPOSAL

TRIPLE RINSE (OR EQUIVALENT), THEN OFFER FOR RECYCLING OR RECONDITIONING, OR DISPOSE OF IN A SANITARY LANDFILL, OR BY INCINERATION IF ALLOWED BY STATE AND LOCAL AUTHORITIES

GENERAL

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES

 **PUREX CORPORATION**
24800 S. MAIN ST./BOX 6200
CARSON, CALIFORNIA 90745

PUREX INSTITUTIONAL

ADMIRAL
ACCEPTED
with COMMENTS
Q in EPA Letter Dated

APR 19 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

1124-102

Disinfectant-Sanitizer
Fungicide-Virucide-Deodorizer
with Organic Soil Tolerance
for Hospital, Institutional

Industrial, School, Dairy and Other Farm Use

BEST AVAILABLE COPY

EPA Reg. No 1124-102

EPA Est. 1124-04-3

Ingredients

Allyl C- 50% C- 40% C- 10% dimethyl	
benzyl ammonium chloride	1.00%
Octyl detyl dimethyl ammonium chloride	2.20%
Didecyl dimethyl ammonium chloride	1.00%
Quaternary dimethyl ammonium chloride	1.00%
Sodium hypochlorite	99.80%

KEEP OUT OF REACH OF CHILDREN.

DANGER: SEE BACK PANEL
FOR FURTHER CAUTIONS.

1 GAL.

It is a violation of Federal law to use this product with its labeling.

Apply ADMIRAL Q with a clean mechanical spray device. Treated surfaces must remain damp until dry or when the use.

DISINFECTION IN HOSPITALS, For disinfecting floors, walls, bedframes, tables, chairs,

Add 3) oz. ADMIRAL Q to 4) surfaces. At this use level, virucidal, Staphylococcus of 55 blood serum when eval.

VIRUCIDAL PERFORMANCE. At in the presence of 55 serum viruses: Herpes simplex 7 on immediate hard non-pore

DISINFECTION IN INSTITUTIONS 4) gallons of water.

FUNGICIDAL PERFORMANCE. At effective fungicide against fungal when used on surface shower and bath areas and using the AOAC Fungicide

DISINFECTION OF BARBER TOOL resors, and scissors) can be of ADMIRAL Q.

DISINFECTION OF POULTRY EQUI waterling founts, feeding or and kernel areas) can be di solution of 2 oz. ADMIRAL Q immersed in this solution, and their feeds must be re troughs, racks and other fo and droppings from floors, poultry or other animals.

After disinfection, ventilate house poultry, or other ani absorbed, set or dried.

All treated equipment that with potable water before

SANITIZING OF FOOD PROCESS LOCATIONS. For sanitizing utensils, dishes, silverware storage and display equipa rinse is required.

Wash and rinse all articles ADMIRAL Q in 4 gallons of for at least one minute fol solution should be prepared For mechanical application applications.

Apply to sink tops, counter and other stationary hard device. No potable water

Dishes, silverware, glassware processing equipment can b solution of ADMIRAL Q. No

At 1 oz./4 gallons ADMIRAL "A" Pasteurized Milk Ordin Services in waters up to 8 by the AOAC Germicide) and and Staphylococcus aureus.

The udders, flanks and tea solution of 1 oz. ADMIRAL is required. Use a fresh solution by dirt and soil. When solution becomes vis