

[1127-3]



# BOURBON

0.75%

**ROTENONE DUST**

RECEIVED

10/3/69

ACTIVE INGREDIENTS—

Rotenone	0.75%
Other Cube Res.	1.15%
INERT INGREDIENT	98.10%

TWO  
POUNDS



NET  
WEIGHT

MANUFACTURED BY  
**THE BOURBON COMPANY**  
 LEXINGTON, KENTUCKY

### CAUTION

Avoid prolonged inhalation of this dust, as it may cause a temporary numbness of the respiratory tract. Breathing fresh air relieves this condition.

Contamination with any body of water may kill fish.

### DIRECTIONS

NOTE: Do not apply to food crops within one day of harvest.

**BEANS:** The Mexican Bean Beetle: This insect feeds from under-side of leaves. Use duster with adjustable nozzle that will direct dust upward, from underneath plant. Use up to 100 lbs. per acre.

**CABBAGE, BROCCOLI:** Cabbage worms, Diamond back Moth. Use up to 66 lbs. per acre.

**PEAS:** Pea Aphids. Use up to 80 lbs. per acre.

Better control can be obtained if plants are covered thoroughly by dusting at the first appearance of insects. Since this dust loses its toxicity in 7 to 10 days after being exposed to sunlight, dustings should be repeated at least every 10 days or more often if necessary. Dusting should be done when foliage is moist with dew.

### FOR SPRAYING:

Add 1 lb. of this dust with about 1 ounce of soap powder to 3 1/3 gallons water. Mix thoroughly and spray plants every 10 days for good control.