

BOURBON

1 3/4% BEAN DUST

CONTAINS SEVIN*



TWO POUNDS MEXICAN BEAN BEETLES &
NET WEIGHT
COLORADO POTATO BEETLES

Active Ingredient:

Carbaryl (1-naphthyl-N-methylcarbamate) 1.75%

Inert Ingredients:

Trade Mark of UCC for 1-naphthyl-N-methylcarbamate



MANUFACTURED BY
THE BOURBON COMPANY
LEXINGTON, KENTUCKY

DIRECTIONS

Apply 1 3/4% BEAN DUST when you first see signs of insect infestation, preferably before feeding begins. For best results apply thoroughly to upper and lower leaf surfaces at the rate of 20 pounds per acre. If needed, repeat applications every 7 to 10 days. Apply just prior to, including the day of, harvest. Do not apply treated beans vines to dairy cattle and hogs.

CROPS

INSECTS

beans

Mexican Bean Beetle

potatoes

Colorado Potato Beetle

CAREFUL WHEN INHALING OR SWALLOWED.
Do not breathe dust and do not take orally. SKIN CONTACT
MAY BE HARMFUL. Avoid skin contact. Wash hands and face
before eating.

NOTE FOR PHYSICIANS: Carbaryl is a moderate irritant.
Cholinesterase inhibitor. Acetanilid is antidote.

AVOID CONTAMINATION OF FOOD.

KEEP OUT OF REACH OF CHILDREN.