DIRECTIONS

Wash Dogs in a solution of 1 part Dip to 100 parts water to destroy Lice and Fleas. Repeat as often as necessary. Wash bedding with 1 part Dip to 50 parts water.

TO CONTROL LIVESTOCK BUILDING CONTAMINATION: Remave all livestock from the building to be disinfected. Thoroughly clean and protect or remove from the building all equipment, including feeders or waterers. Remove all litter or bedding and clean the building thoroughly.

Apply disinfectant at the recommended rate of 1 cup to a 10-quart pail of water. When feeding and watering equipment are disinfected thoroughly, scrub with detergent and rinse with potable water. Ventilate the building and do not house poultry or livestock until treatment has been absorbed, set or dried.

Do not make any disinfectant applications when animals are in the building.

Strict adherence to regular scheduled cleaning and disinfecting program aids in reducing contamination in animal quarters.

Do not use in milk rooms or milking parlors.

WARNING: KEEP OUT OF REACH OF CHILDREN.

May produce severe burns. May be fatal if swallowed or absorbed through skin. If swallowed, give white of egg, mucilage or starch paste an antidote. Use 1 tablespoon full of mustard in glass of water as an emetic. Call a physician.

Do not get in eyes, on skin or on clothing. Wash thoroughly with water after handling. Avoid breathing vapor fumes or spray mist. Keep away from domestic animals. Avoid contamination of feed and foodstuffs. Discard empty container after repeated rinsing with water.


CAUTION: No not use on poultry. Do not use in poultry houses. DO NOT USE ON CATS.



NET CONTENTS ONE C
BOURBON
AND
DISINFECT
Active Ingredients: Coal Tar Hydrocarbons Soap Coal Tar Acids Inert Ingredients: Water PHENOL COEFFICIENT 5 — A.O.A.O
An Efficient Insecticide, Deoc Cleanser
DOGS
 LIVESTOCK BUILD HOME PREMISES
FOR EXTERNAL USE
Distributed by THE BOURBON COM Manufacturers and Distribute LEXINGTON, KENTUC
WARNING: Keep out of read
See Antidote statement an

See Antidote statement and other required warning statements on the back panel.

1 1 1





.C. METHOD dorizer,

DINGS



IPANY ors KY.

ch of children.



•

DIRECTIONS FOR HOME USE

FOR DISINFECTION AGAINST SALMONELLA CHOLERAESUIS:

Use one-half cup of Coal Tar Disinfectant to each 10quart pail of water taken for mopping, scrubbing, sprinkling or cleaning purposes.

GARBAGE: To deodorize thoroughly moisten the gar-bage with a strong solution of this product. Rinse the empty can with some of the solution and leave the wash in the can. Coal Tar Disinfectant helps to prevent the breeding of flies in the container.

FLOORS: Use one-half cup of Coal Tar Disinfectant to a 10 quart pail of water when scrubbing or mopping.

KITCHEN SINKS: Scrub above and beneath sink, along the sides and edges of the sink, using a solution of one-half cup of Coal Tar Disinfectant to a 10 quart pail of water, and flush pipes with the remained of the fluid.

WATER CLOSETS AND URINALS: Wash the closet and urinal bowls regularly with Coal Tar Disinfectant solution in the proportion of one-half cup to a gallon of water. Scrub the floors and seat frequently with Coal Tar Disinfectant Solution.

USDA Reg. No. 1127-8

H 7-71

WITH COMMENTS ACCEPTED

7

ACC

NITE

AS A DISINFECTANT for use against the germs of typhoid fever and hog cholera on environmental surfaces use a solution of one part Bourbon Pine Oil Disinfectant to 100 parts water or one tablespoonful cf disinfectant to three pints of water. Feeding utensils to be disinfected should be exposed to the action of this solution for at least five minutes and rinsed with potable water prior to reuse.

FOR DISINFECTING AND DEODORIZING floors and walls use a solution of one part Bourbon Pine Oil Disinfectant to 50 parts water, or two ounces of disinfectant to each gailon of water to mop or wipe the immediate area.

IN GARAGE RECEPTACLES --- To aid in checking the development of putrefactive action and the breeding of flies remove all gross filth and spray freely with a solution of one part disinfectant to 50 parts water.

filth and trash.

IN POULTRY HOUSES --- To disinfect poultry houses use a solution of one part disinfectant to 50 parts water. Remove all poultry and feeds from premises, trucks, coops, and crates. Empty all troughs, racks, and other feeding and water appliances. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry. Saturate surfaces with solution and ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried. All treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers must be thoroughly rinsed with potable water prior to reuse.

TO CLEAN PAINT BRUSHES and to keep them from hardening use a solution of 1 part disinfectant to 5 parts water.

If swallowed do not induce vomiting. Drink a large quantity of milk, egg whites or gelatin solution. If these are not available drink large quantities of water, avoid alcohol. Call a physician.

DO NOT USE AROUND THE HOUSE OR RECREATION AREAS. RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD IT.

DIRECTIONS FOR USING


BOURBON PINE OIL DISINFECTANT

IN DOG KENNELS - To kill fleas spray the premises thoroughly with a solution of one part disinfectant to 50 parts water after removing all

CAUTION

HARMFUL IF SWALLOWED. CONCENTRATE MAY CAUSE IRRITA-TION. AVOID CONTACT WITH SKIN AND EYES.

EPA Reg. No. 1127-63



A disinfectant and deodorant for inanimate objects.

THE BOURBON COMPANY