[144-]

CAUTION COMBUSTIELE MIXTURE Permit Ne. M133036

KILLS Red Mites - Fleas - Flies - Flying Moths - Reaches - Mesquitoes Ticks and Red Mites on Caged Birds When Used EXTERNALLY AND AS DIRECTED.

To kill roaches, water bugs and silver fish: spray a heavy mist of "Mit-E-Kil" inside cracks and crevices around sinks, pipes and drains where these insects hide when they emerge spray again.

For gnats, flies, flying moths & mosquitoes: Close doors and windows and spray toward ceiling until room is filled with heavy mist. Leave closed for 10 minutes. Repeat application as often as necessary.

DO NOT REMAIN IN TREATED AREAS AND VENTILATE THE AREA AFTER TREATMENT IS COMPLETED.

۰.

Mit-E-Kil

THE NEW IMPROVED FORM OF

Insect Control

for Fleas on dogs & cats and Red Mites on birds

100% ACTIVE INGREDIENTS

DO NOT SPRAY NEAR OPEN FLAME

HARMFUL IF SWALLOWED

CAUTION KEEP OUT OF REACH OF CHILDREN

U.S.D.A. Bog. No. 1144-1

Active Ingredients 100%

Pyrethrins _____ *Technical Piperonyl Butaxide ______ 1,890% Petroloum Hydrocarbons _____ 97,735% *equivalent to 1.512% (Butylcarbityl) (6-propylpiperonyl) ether and 0.376% related compounds.

NET 32 FL. OZ. (1 QT.)

HINTON & CO. INC. P. O. BOX 26 FARMINGDALE, N. Y. 11735

1144-1



.375%

PIRECTIONS

For Figas on Bers;

Hold dog with one hand and spray lightly. Also spray badding and other items with which does come in contact. When spraying Lennel, deghouse or room, spray entire space and spray toward ceiling until complete area is filled with heavy mist.

For ticks. Brush back dogs hair and spray lightly as -Mit-E-Kill' will kill ticks only on contact. Repeat application as often as necessary.

For Fleas an Cats:

Hold cat with one hand and spray lightly. Also soray bedding and all other items with which cat comes in contact. Repeat application as often as necessary.

For Red Mites on Birds. **Pigeons and Chickens:**

Close all doors and windows. Spray birds, cages and entire room including cracks and floor. Spray toward ceiling and until room is filled with light mist. Spray as often as necessary.

For Canaries Hold sprayer at least two feet away from the cages and spray directly into cages Spray as often as necessary.

DO NOT REMAIN IN TREATED AREAS AND VENTILATE THE AREA AFTER TREATMENT IS COMPLETED.

- - • •