

## CAUTION

Causes irritation. Avoid contact with eyes, skin, and clothing. Contains petroleum solvent. Avoid inhalation of spray mists. Avoid contamination of foodstuffs.

Keep out of reach of children. If swallowed, call a physician immediately. In case of eye contact, flush with large amount of water for 15 minutes; get prompt medical attention. Wash thoroughly after handling.

## OBSERVE THESE PRECAUTIONS

Do not use in or near areas and the home, recreation areas, or similar sites. Avoid application of spray or drift to spray to desirable plants, as this product may be toxic to certain garden and tree vegetables, flowers, grapes, fruit trees, and ornamentals, other desirable plants. Even minute quantities of this spray may harm evergreen trees. Avoid application made by airplane over the type of crops or sites listed to avoid injury where there is no hazard from drift. All sprays are likely to drift under certain conditions. The lower the spray, the greater the likelihood of drift effect. Coarse sprays are more likely to drift.

At water temperatures up to 120° F., may cause injury to susceptible plants grown in trays.

Do not use on freshly seeded pastures until grass has become well established. Most grasses are usually 14 days old or older.

Do not use in areas that contain water dependent fish or other aquatic life. Excess amounts of this product in the water may temporarily kill or retard growth of the aquatic life.

Do not use in areas where there are susceptible plants or animals growing.

Respect the toxicity of this herbicide. Do not use in areas where it is prohibited. Do not use in areas where it is restricted. Do not use in areas where it is prohibited.

Do not use in areas where it is prohibited. Do not use in areas where it is restricted. Do not use in areas where it is prohibited.

Do not use in areas where it is prohibited. Do not use in areas where it is restricted. Do not use in areas where it is prohibited. Do not use in areas where it is restricted. Do not use in areas where it is prohibited.

Do not use in areas where it is prohibited. Do not use in areas where it is restricted. Do not use in areas where it is prohibited. Do not use in areas where it is restricted. Do not use in areas where it is prohibited.

Do not use in areas where it is prohibited. Do not use in areas where it is restricted. Do not use in areas where it is prohibited. Do not use in areas where it is restricted. Do not use in areas where it is prohibited.

## NON-WARRANTY NOTICE

Read and follow directions carefully. Seller warrants such directions to be proper and that the product conforms to the descriptions on this label. Timing and method of application, weather and crop conditions, and other influencing factors are beyond the control of the Seller, so Seller makes no other warranty, express or implied, including a warranty of merchantability, with reference to the results or effect of use. No person has authority to make a representation in regard to this label.

## THIS CONTAINER IS NON-RETURNABLE

REG. NO. 114-114-114  
0-51



# Amoco

## 2,4,5-T LV ester

### FOR BRUSH CONTROL


CAUTION: KEEP OUT OF THE REACH OF CHILDREN. SEE OTHER CAUTION STATEMENTS ON SIDE PANEL.

A selective, emulsifiable or oil dilutable herbicide for the control of most broadleaved annual weeds and many herbaceous and woody perennials. This product contains the powerful low volatile ester of 2,4,5-T which will control certain plants more efficiently than with 2,4-D. It is effective in controlling practically all weedy and brushy species found in pastures, fence rows, and along highways, and railroads, and in right-of-ways for power, telephone, and pipe lines—including Alder, Aspen, Birch, Brambles (Raspberry and Blackberry), Chokecherry, Elder, Elderberry, Elm, Hawthorn, Hickory, Honeysuckle, Locust, Oaks, Osage Orange, Poison Ivy, Poison Oak, Salmon Berry, Sassafras, Sumac, Sweet Gum, Wild Cherry, Wild Grape, Wild Rose, Willow, and many others.

## FOLLOW DIRECTIONS CAREFULLY

### To prepare a spray

Always mix with water or oil (AMOCO® Diesel Fuel, AMERICAN® Heater Oil, or AMERICAN® Kerosene) so as to apply the recommended amount of AMOCO 2,4,5-T LV Ester in the amount of diluent necessary to cover the given



area with the spraying equipment used. Add half of the required amount of water or oil to a spray tank, then add AMOCO 2,4,5-T LV Ester with agitation, and finally the balance of water or oil, with continued agitation. **Warning:** If AMOCO 2,4,5-T LV Ester is to be used in preparing straight oil mixtures, do not let water get into the AMOCO 2,4,5-T LV Ester itself or into the finished mixture. **Note:** AMOCO 2,4,5-T LV Ester forms an emulsion with water, which tends to separate on standing for prolonged periods of time. Provide agitation to prevent such separation and to insure uniformity of spray mixture.

**For mixing small quantities of spray,** one tablespoonful of AMOCO 2,4,5-T LV Ester in 1½ gallons of oil or water is approximately equivalent to 1 quart in 100 gallons.

## AMOUNTS OF SPRAY MIXTURE TO APPLY

**Airplane Application - Water:** Dilute recommended amount of AMOCO 2,4,5-T LV Ester in 3 to 5 gallons of water and apply per acre.

**Oil:** Dilute recommended amount in 1 gallon of AMERICAN Heater Oil or AMOCO Diesel Fuel and apply per acre.

PRODUCT OF AMERICAN OIL COMPANY  
CHICAGO, ILLINOIS, USA 60680

**Ground Sprayers:** Dilute recommended amount in 8 to 20 or more gallons of water and apply per acre. The amount of water required will depend upon the crop and type of equipment used. Thorough agitation of spray mixture is necessary for best results.

## FOLIAGE TREATMENT

Spray woody growth up to 6 or 8 feet tall after foliage is well developed using a drenching spray containing 3 to 4 quarts of AMOCO 2,4,5-T LV Ester per 100 gallons of water. Taller brush can be sprayed successfully, although in many cases basal bark or stump treatment is preferable. Coverage should be complete, and all parts of plants, including foliage, stem, stems and bark, should be wet with spray. Best results usually will be obtained from applications made soon after maximum foliage development in the spring. Later applications may be made if good growing conditions prevail. Power equipment with pressures up to 250 pounds per square inch obtaining satisfactory spray coverage. Repeat applications may be required as new growth appears. Usually, a single treatment in a year is usually sufficient. Do not apply by airplane in the vicinity of cotton or other 2,4,5-T susceptible crops. Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

For treatment of trees and brush in pastures apply AMOCO 2,4,5-T LV Ester at rate of 3 quarts per acre in sufficient water or oil to give adequate coverage of foliage depending on whether application is made by ground sprayer or airplane.

## BASAL BARK TREATMENT

Brush and small trees can be controlled by spraying the basal parts of brush and tree trunks to a height of 12 to 15 inches from the ground line. Use a solution of 1 gallon of AMOCO 2,4,5-T LV Ester in 24 gallons of oil, mix thoroughly. With certain easier to control species, 3 quarts of AMOCO 2,4,5-T LV Ester in 24 gallons of oil is effective. Knapsack or power equipment may be used, but complete wetting of the indicated area is necessary, particularly at the ground line. This means spraying until run-down or run-off to the ground line is noticeable. Old or rough bark requires more spray volume than young or smooth bark. Low pressures are desirable. Apply at any time, including the winter months. Delayed response and killing can often be expected, treated brush or trees therefore should not be cut for a period of one year following application.

## STUMP TREATMENT

When growth is more than 6 to 8 feet tall, cut it close to the ground and spray the stumps and stubs. Use a thoroughly mixed solution of 1 gallon of AMOCO 2,4,5-T LV Ester in 24 gallons of oil. All exposed bark, as well as cut surfaces should be wet thoroughly. This means spraying until run-down or run-off to the ground line is noticeable. Old or rough bark requires more spray volume than young or smooth bark. Apply at any time, including the winter months, preferably to freshly-cut stumps. Best results are usually obtained on stumps two inches across or larger.

## "FRILL" TREATMENT

For large trees, make a single-hack girdle or "frill" of overlapping axe cuts at a convenient height. Treat the injured area with a mixture of 3 to 4 quarts of AMOCO 2,4,5-T LV Ester in 24 gallons of oil.

## INJECTION TREATMENT

Make injections spaced 2 inches apart (edge to edge) as near the root collar as possible. In the case of resistant species such as Hickory, injections should overlap around the tree. Mix 1 gallon of AMOCO 2,4,5-T LV Ester with 20 gallons of oil and apply 5 cubic centimeters of the solution per injection. Injections may be made anytime during the year, however they are most effective when made in the spring or early summer.

MADE IN U.S.A. 744 (D) MP 1-6571

# CONTENTS      GALLONS



# Amoco

## 2,4,5-T LV ester

**FOR BRUSH CONTROL**

**CAUTION: KEEP OUT OF THE REACH  
OF CHILDREN. SEE OTHER CAUTION  
STATEMENTS ON SIDE PANEL.**

<b>ACTIVE INGREDIENTS BY WEIGHT</b>	
Isocetyl ester of 2,4,5-Trichlorophenoxyacetic acid	66.0%
(Equivalent of 45.1% 2,4,5-Trichlorophenoxyacetic acid)	
<b>INERT INGREDIENTS BY WEIGHT</b>	<b>34.0%</b>
<b>TOTAL</b>	<b>100.0%</b>

1 gal. of AMOCO 2,4,5-T LV Ester contains the equivalent  
of 4 pounds of 2,4,5-Trichlorophenoxyacetic Acid