

DIRECTIONS

AMOCO Ronnel Spray mixes readily with water or oil. To prepare a finished spray, see chart below.

Type and Use	% Ronnel	Amount Ronnel Spray to use for			Remarks
		1 gal.	3 gal.	25 gal.	
LIVESTOCK INSECTS —Repeat as necessary but not oftener than every 2 wks. Cattle and Calves: Hornflies and Lice. Lone Star and Winter Ticks Sheep, Goats, Lambs and Kids: Lice, Sheep Keds, Wool Maggots and Hornflies	½% ¾% ½%	5 tbs. 7½ tbs. 5 tbs.	½ pt. ¾ pt. ½ pt.	2 qt. 3 qt. 2 qt.	Dilute with water only. Spray entire animal to wet hair coat thoroughly.
Cattle Backrubber— For Hornflies. Place where animals travel regularly or congregate.					Mix 1 gallon AMOCO Ronnel Spray with 21 gallons Diesel or similar oil. Recharge as necessary. Do not mix with used motor oil.
FLY BAIT— Where Flies congregate such as floors, windows, doorways.	2%	¾ pt.	1 qt.	2 gal.	Add 3 ounces of sugar per gallon diluted spray. Spray or sprinkle at 1 to 2 day intervals until control is obtained.
COCKROACH CONTROL— Use 1% spray for German Roaches, 2% for others.					Apply as a spot spray to all areas frequented by Cockroaches.
Buildings— Including dwellings, institutions and food handling establishments.	1% or 2%	½ pt. or ¾ pt.	1 pt. or 1 qt.	1 gal. or 2 gal.	
Restaurants, Supermarkets— Oil mist spray inside buildings for German Cockroaches.	1% or 2%	½ pt. or ¾ pt.	1 pt. or 1 qt.	1 gal. or 2 gal.	Dilute with oil and apply by remote control when buildings are not occupied. Cover exposed food and equipment. Use 1 gallon diluted spray per 50,000 cu. ft.
Food Processing Plants	1% or 2%	½ pt. or ¾ pt.	1 pt. or 1 qt.	1 gal. or 2 gal.	Spray when not in operation. Cover produce.
Meat Packing Plants	2%	¾ pt.	1 qt.	2 gal.	Spray outer and inedible product areas.
Poultry Processing Plants	2%	¾ pt.	1 qt.	2 gal.	Spray outer and inedible product areas.

U. S. PATENT No 2,599,516

USDA REG. NO. 1145-30 (028)

MADE IN U. S. A. 754 - OX - MP - L - 1068

ACCEPTED

AMOCO Ronnel Spray



PRODUCT OF AMERICAN OIL COMPANY
CHICAGO, ILL. USA 60680

CAUTION KEEP OUT OF REACH OF CHILDREN SEE OTHER CAUTION STATEMENTS ON LABEL

LIVESTOCK REMEDY, NOT FOR HUMAN USE

A water emulsifiable, oil soluble, insecticide concentrate for Residual Fly Control in farm buildings, and control of other livestock pests, Roaches, Flies.

FOR LIVESTOCK, FARM, RANCH, AND INDUSTRIAL INSECT CONTROL

OBSERVE THESE PRECAUTIONS

Do not spray beef cattle within 8 weeks, sheep and goats within 12 weeks of slaughter. Do not spray lactating dairy animals, or dry dairy animals within 21 days of the lactating period. Backrubbers may be used for beef animals to within two weeks of slaughter, on dry dairy cows to within three weeks of freshening. Not for use on lactating dairy cows. Do not use in rooms where milk is processed or stored. Do not allow spray to come in contact with food, feed, or water. Remove lactating dairy animals from buildings during application of residual fly spray. Do not use on livestock receiving an organic phosphate treatment from any other source at the time of treatment.

NOTE TO PHYSICIAN: This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ACTIVE INGREDIENTS (by weight)

Ronnel® 0,0-dimethyl 0-(2,4,5-trichlorophenyl) phosphorothioate	23.7%
Aromatic petroleum derivative solvent	59.1%
INERT INGREDIENTS (by weight)	17.2%
TOTAL	100.0%

*From technical grade not suitable for internal use in animals.

1 gallon AMOCO Ronnel Spray contains the equivalent of 2.0 pounds Ronnel

CAUTION

INJURIOUS TO THE EYES, IRRITATING TO THE SKIN, HARMFUL IF SWALLOWED

Do not get in the eyes. Avoid prolonged or repeated contact with the skin. If swallowed, call a physician immediately. Unless instructed otherwise, induce vomiting by introducing a finger into throat or by giving warm salt water (one tablespoon of salt to a glass of warm water). Avoid prolonged breathing of spray mists. Avoid contamination of food, feedstuffs, and water supplies. Keep container closed. Wash thoroughly with soap and water after handling. Do not dispose of any unused material in such a manner that contact with humans or beneficial animals may occur.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated.

COMBUSTIBLE: Keep container closed and away from heat and open flame.

DIRECTIONS

Type and Use	% Ronnel	Amount Ronnel Spray to use for			Remarks
		1 gal.	3 gal.	25 gal.	
RESIDUAL FLY SPRAY— A ½% spray will give about 2 weeks control. A 1% spray will give 4-6 weeks control.					Spray walls, ceilings, partitions to obtain complete coverage with no run-off. Use 1 gallon of diluted spray per 500-1000 sq. ft.
Farm Buildings— Including poultry houses, livestock shelters, dairy and hog barns.	½% or 1%	5 tbs. or ½ pt.	½ pt. or 1 pt.	2 qt. or 1 gal.	Add 3 ounces of sugar per gallon for added effectiveness. Dilute with water only.
Food Processing Plants	½% or 1%	5 tbs. or ½ pt.	½ pt. or 1 pt.	2 qt. or 1 gal.	Spray when not in operation. Cover produce and equipment. Spraying outside area will improve control.
Dwellings— Outside areas, including porches, patios, garages.	½% or 1%	5 tbs. or ½ pt.	½ pt. or 1 pt.	2 qt. or 1 gal.	Spray to point of run-off.
Restaurants, Supermarkets— Outside area.	½% or 1%	5 tbs. or ½ pt.	½ pt. or 1 pt.	2 qt. or 1 gal.	Spray outside walls, receiving docks, doorways, and refuse areas.
Meat Packing Plants	1%	¾ pt.	1 pt.	1 gal.	Spray outer and inedible product areas.
Poultry Processing Plants	1%	¾ pt.	1 pt.	1 gal.	Spray outer and inedible product areas.
Refuse areas and containers	1%	¾ pt.	1 pt.	1 gal.	Spray to soak all exposed surfaces.
Caged Layers on wire.	2%	¾ pt.	1 qt.	2 gal.	Spray poultry droppings beneath cages.
MISCELLANEOUS INSECTS					Apply 1 gal. of dilute spray per 1000 sq. ft.
Fleas and Spiders in buildings, laundry trucks, dog kennels.	1%	¾ pt.	1 pt.	1 gal.	Spray floors, walls, cracks, crevices, sleeping quarters of pets, where insects breed and harbor.
Bedbugs in buildings and sleeping quarters, on beds, and bedding.	1%	¾ pt.	1 pt.	1 gal.	Spray mattress and bedding tightly, including seams and folds, cracks and crevices in bed frames, walls, baseboards, and floors.
Fleas in lawns and kennel runs.	1%	¾ pt.	1 pt.	1 gal.	Spray infested areas. Keep children off treated area until spray has dried.
Ants, Mosquitoes and Silverfish	1%	¾ pt.	1 pt.	1 gal.	Spray areas where insects frequent.

NON-WARRANTY NOTICE

Follow directions carefully. Seller warrants such directions to be proper and that product conforms to the descriptions on this label. Timing and method of application, weather and crop conditions, and other influencing factors in the use of the product are beyond the control of the Seller, so Seller makes no other warranty, express or implied (including a warranty of merchantability), with reference to the results or effects of use. No person has authority to make any representation not contained on this label.

Net Contents

U.S. GALS.