

[1145-34]

CAUTION

Avoid contact with skin. Harmful if swallowed. If swallowed, call a physician immediately. Avoid contamination of milk and milk handling equipment. Avoid overspraying which will result in wetting animal's hide. Use according to directions.

COMBUSTIBLE: Do not use or store near heat or open flame.

NON-WARRANTY NOTICE

Follow directions carefully. Seller warrants such directions to be proper and that product conforms to the descriptions on this label. Timing and method of application, weather and crop conditions, and other influencing factors in the use of the product are beyond the control of the Seller, so Seller makes no other warranty, express or implied (including a warranty of merchantability), with reference to the results or effects of use. No person has authority to make any representation not contained on this label.

USDA REG. NO. 1145-34
2671


BOVINOL[®] SUPER Stock Spray

Specially formulated for use in barn foggers



PRODUCT OF **AMERICAN OIL COMPANY**
CHICAGO, ILL. USA 60686

CAUTION KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTION STATEMENTS ON SIDE PANEL



ACTIVE INGREDIENTS (by weight)	
Pyrethrins	0.10%
Piperonyl butoxide, technical*	0.20%
N-octyl bicycloheptene dicarboximide	0.33%
Di-n-propyl isocinchomeronate**	0.20%
Petroleum distillate	99.17%
TOTAL	100.00%

*Equivalent to 0.16% butylcarbitol 6-propylpiperonyl ether and 0.04% of related compounds
**MGM Repellent R-324

READY TO USE

DIRECTIONS FOR USE

FOR BARN FOGGING: To control House Flies, Stable Flies, Horn Flies, Mosquitoes, and Gnats, use approximately one ounce of BOVINOL Super Stock Spray for each 1000 cubic feet of space. Have cattle in barn before spraying. Keep doors and windows closed for at least 10 minutes. Sweep up and destroy all fallen flies. Follow recommendations of equipment manufacturer for proper positioning of the fogging machine.

FOR DAIRY CATTLE SPRAYING: To protect dairy stock from biting Flies, House Flies, Mosquitoes, and Gnats, as well as certain other flying pests, when barn foggers are not used, spray daily before milking, preferably in the barn to avoid wastage due to wind drift, using one ounce per head. Larger quantities may cause skin irritation or inhibit skin ventilation in hot weather. Spray lightly, but evenly and thoroughly over every part of the animal, with particular attention to the back, flanks, and legs—avoid udder and area around eyes. Do not soak the skin or wet the hide. Do not apply on calves. Do not blanket, brush, or curry the animals shortly after spraying, as this might increase the risk of skin irritation.

FOR HORSES: Spray lightly or sponge horses, using one ounce per adult animal, proportionately less for smaller animals. All parts of the animal's body should be covered, with special attention to shanks, legs, and shoulders. Apply till hair is moistened, but do not soak hide. Do not get the spray under any leather, such as bridle, harness, or saddle, as the rubbing under leather is apt to produce irritation. Also, do not blanket, brush, or curry the animals shortly after spraying.

MADE IN U.S.A.

Net Contents

U.S. GALS.