



AMOCO[®] Granular 2,4-D For Corn

20/35 Mesh Size



PRODUCT OF **AMERICAN OIL COMPANY**
CHICAGO • ILL. • USA

CAUTION KEEP OUT OF REACH OF CHILDREN • SEE OTHER CAUTION STATEMENTS ON LABEL

ACTIVE INGREDIENTS (by weight)
Isooctyl ester of 2,4-Dichlorophenoxyacetic acid 31.08%

A 2,4-D low volatile ester granular weed killer
for control of certain weeds in corn

CAUTION KEEP OUT OF REACH OF CHILDREN · SEE OTHER CAUTION STATEMENTS ON LABEL

| | |
|---|----------------|
| ACTIVE INGREDIENTS (by weight) | |
| Isooctyl ester of 2,4-Dichlorophenoxyacetic acid . . . | 31.08% |
| (Equivalent to 20.0% 2,4-Dichlorophenoxyacetic acid) | |
| INERT INGREDIENTS (by weight) | 68.92% |
| TOTAL | <u>100.00%</u> |

CAUTION

Do not swallow. Avoid contact with skin, eyes, and clothing. Avoid inhaling dust. May cause skin irritation. Avoid contamination of foodstuffs.

OBSERVE THESE PRECAUTIONS

Never use AMOCO Granular 2,4-D except as recommended on this label. Never treat, or allow particles or dust to drift on desirable crops or plants, as this product, even in minute quantities, may cause severe injury to cotton, vegetables, ornamentals, flowers, grapes, tomatoes, soybeans, tobacco, fruit trees, and other plants. Do not apply on a windy day. Use amounts recommended—do not overdose. Although AMOCO Granular 2,4-D is formulated with a low volatile ester of 2,4-D, in order to reduce the hazard of vapor damage, every precaution must be exercised to prevent injury to nearby susceptible plants.

Do not store or place near fertilizers, seed, plants, insecticides, or fungicides. Do not re-use empty container. Destroy by burning in a safe place. Do not use in greenhouses. Avoid contamination of irrigation water and reservoirs or other areas used for drinking water supplies.

It is safest to use separate equipment for applying AMOCO Granular 2,4-D. Clean thoroughly if equipment is used for other purposes. Do not contaminate water used for spray purposes.

A 2,4-D low volatile ester granular weed killer for control of certain weeds in corn (pre-emergence treatment only).

SPECIFIC USE RECOMMENDATIONS

CORN (Midwest only)—To control germinating broad-leaved weeds in corn such as Cocklebur, Lambsquarter, Mustard, Morning Glory, Pigweed, Purslane, and Ragweed apply AMOCO Granular 2,4-D at planting time or before either the crop or the weeds appear above the soil surface. **DO NOT USE ON LIGHT OR SANDY SOILS.** Do not disturb soil surface after applying granules—allow granules to remain on soil surface. Dry weather after application will reduce the effectiveness of the treatment. Heavy rains after treatment may cause some injury to germinating corn seed and result in some early crop stunting. This risk may be reduced by planting the corn one inch deeper than normal (2½—3 inches deep).

For overall or broadcast treatments, apply 10 pounds of AMOCO Granular 2,4-D per acre, using broadcast applicators or fertilizer spreaders that can apply small amounts of granules evenly. For band applications, use 3¼ pounds per acre (13 inch band with 40 inch corn rows). Use corresponding rates for other row spacings. Use granular applicator for uniform distribution.

NON-WARRANTY NOTICE

Follow directions carefully. Seller warrants such directions to be proper and that product conforms to the descriptions on this label. Timing and method of application, weather and crop conditions, and other influencing factors in the use of the product are beyond the control of the Seller, so Seller makes no other warranty, express or implied (including a warranty of merchantability), with reference to the results or effects of use. No person has authority to make any representation not contained on this label.

KEEP BAG CLOSED WHEN NOT IN USE