

## CAUTION

Causes irritation. Avoid contact with eyes, skin, and clothing. Avoid inhalation of spray mists. Avoid contaminating of foodstuffs.

Harmful if swallowed. If swallowed, call a physician immediately. In case of eye contact, flush with large amounts of water for 15 minutes, get prompt medical attention. Wash thoroughly after handling.

## OBSERVE THESE PRECAUTIONS

Do not apply AMOCO MCPA Weed Killer-M directly to, or otherwise permit it to come into contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton, or other desirable plants which are sensitive to MCPA and similar materials, and do not permit spray mist containing it to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. (Coarse sprays are less likely to drift.) Applications by airplane, ground rigs, and hand dispensers should be carried out only when there is no hazard from drift.

Do not apply by airplane in the vicinity of cotton, grapes, or other desirable MCPA susceptible crops.

Do not contaminate irrigation ditches or water used for domestic purposes.

Do not store near fertilizers, seeds, insecticides, or fungicides. Excessive amounts of 2-methyl-4-chlorophenoxyacetic acid in the soil may temporarily inhibit seed germination or plant growth.

To avoid injury to desirable plants, do not store, handle, or apply other agricultural chemicals with the same containers or equipment used for AMOCO MCPA Weed Killer-M.

Local conditions may affect the use of herbicides. Extension Service Weed Specialists or Agricultural Experiment Stations in many states issue recommendations to fit local conditions.

Be sure that use of this product conforms to all applicable regulations.

For timing and specific dosage recommendations consult your local Extension Service Weed Specialist or Agricultural Experiment Station.

Obtain necessary approval and/or permits for use in States or areas where required.

AMOCO MCPA Weed Killer-M exposed to sub-freezing temperatures should be warmed to at least 40°F. and mixed thoroughly before using.

Do not re-use empty container. Rinse equipment and containers and dispose of wastes by burying in non-crop areas away from water supplies. Dispose of containers by punching holes in them and burying with wastes.

## NON-WARRANTY NOTICE

Follow directions carefully. Seller warrants such directions to be proper and that product conforms to the descriptions on this label. Timing and method of application, weather and crop conditions, and other influencing factors in the use of the product are beyond the control of the Seller, so Seller makes no other warranty, express or implied (including a warranty of merchantability), with reference to the results or effects of use. No person has authority to make any representation not contained on this label.

**THIS DRUM IS NON-RETURNABLE - NO REFUND**



# Amoco

# MCPA weed killer

**CAUTION: HARMFUL IF SWALLOWED**  
KEEP OUT OF THE REACH OF CHILDREN  
SEE OTHER CAUTION STATEMENTS ON  
SIDE PANEL

MFD FOR AMERICAN OIL COMPANY  
CHICAGO, ILL. USA 60680

**NET GALLONS**

ACTIVE  
Dimeth  
4 Chlor  
12 Meth  
equival  
INERT  
TOTAL  
1 gallon  
contain  
2 Meth

**GR**

ions may affect the use of herbicides  
 Service Weed Specialists or Agricultural  
 Stations in many states issue recommen-  
 local conditions.

use of this product conforms to all  
 gulations

id specific dosage recommendations con-  
 al Extension Service Weed Specialist or  
 Experiment Station.

sary approval and/or permits for use in  
 as where required.

CPA Weed Killer-M exposed to sub-freez-  
 ures should be warmed to at least 40°F.  
 thoroughly before using.

empty container. Rinse equipment and  
 nd dispose of wastes by burying in non-  
 rway from water supplies. Dispose of  
 y punching holes in them and burying

**WARRANTY NOTICE**

ions carefully. Seller warrants such direc-  
 roper and that product conforms to the  
 on this label. Timing and method of ap-  
 eather and crop conditions, and other  
 actors in the use of the product are be-  
 ontrol of the Seller, so Seller makes no  
 ty, express or implied (including a war-  
 chantability), with reference to the results  
 use. No person has authority to make  
 ation not contained on this label!

**THIS IS NON-RETURNABLE - NO REFUND**



**Amoco**®

**MCPA  
 weed killer-M**

**CAUTION: HARMFUL IF SWALLOWED  
 KEEP OUT OF THE REACH OF CHILDREN  
 SEE OTHER CAUTION STATEMENTS ON  
 SIDE PANEL**

**MFD FOR AMERICAN OIL COMPANY  
 CHICAGO, ILL USA 60680**

**NET GALLONS**

**ACTIVE INGREDIENTS BY WEIGHT**  
 Dimethylamine salt of 2 Methyl  
 4 Chlorophenoxyacetic acid MCPA 52.1%  
 (2 Methyl 4 Chlorophenoxyacetic acid  
 equivalent 47.5%)  
**INERT INGREDIENTS BY WEIGHT** 47.9%  
**TOTAL** 100.0%  
 1 gallon of AMOCO MCPA Weed Killer M  
 contains the equivalent of 4 pounds of  
 2 Methyl 4 Chlorophenoxyacetic acid

**GROSS POUNDS**

**FOLLOW DIRECTIONS CAREFULLY**

In preparing spray mixtures add at least half the  
 water to the spray tank before adding the AMOCO  
 MCPA Weed Killer-M. Apply at moderate pressure by  
 air or ground equipment during warm weather when  
 weeds are growing actively.

**WEED CONTROL  
 IN FLAX**

For control of Mustard and other susceptible weeds  
 use 1/2 pint of AMOCO MCPA Weed Killer-M per acre  
 in the necessary amount of water for uniform dis-  
 tribution. Treat flax when 3 to 6 inches tall before  
 buds begin to form.

**WEED CONTROL  
 IN SMALL GRAINS  
 (OATS, BARLEY, AND WHEAT)**

For control of Mustard, Lambsquarters, and other sus-  
 ceptible weeds use 1/2 pint AMOCO MCPA Weed  
 Killer M in 5 to 10 gallons of water per acre as soon  
 as possible after weeds emerge but not before seed-  
 ling grain is 4 inches tall and together with weed  
 seedlings they form a canopy over clover seedlings.  
 Treatment may then be made up until grain is 8 to 10  
 inches tall (in tiller to early boot stage). With certain  
 weed species that are more difficult to control and  
 under conditions where normally easy to control

species are more resistant to weed  
 3/4 to 1 pint of AMOCO MCPA We  
 required for effective weed cont  
 small grain interplanted with swee  
 trefoil unless injury to the legum  
 Use on grain interplanted with s  
 where specifically recommended  
 Service Weed Specialist or Agri  
 Station. MCPA is safer on oats th

**WEED CONTROL  
 IN ESTABLISHED  
 ALFALFA AND RED CLOVER**

For control of Yellow Rocket and  
 winter annual weeds use 1 pint  
 Weed Killer-M in the amount of  
 good coverage. Application mus  
 early spring before growth of the  
 commenced. Temperature at time  
 be above 40°F.

MCPA REG. NO. 1145-127  
 1971

## FOLLOW DIRECTIONS CAREFULLY

In preparing spray mixtures add at least half the water to the spray tank before adding the AMOCO MCPA Weed Killer-M. Apply at moderate pressure by air or ground equipment during warm weather when weeds are growing actively.

### WEED CONTROL IN FLAX

For control of Mustard and other susceptible weeds, use 1/2 pint of AMOCO MCPA Weed Killer-M per acre in the necessary amount of water for uniform distribution. Treat flax when 3 to 6 inches tall before buds begin to form.

### WEED CONTROL IN SMALL GRAINS (OATS, BARLEY, AND WHEAT)

For control of Mustard, Lambsquarters, and other susceptible weeds use 1/2 pint AMOCO MCPA Weed Killer M in 5 to 10 gallons of water per acre as soon as possible after weeds emerge, but not before seedling grain is 4 inches tall and together with weed seedlings they form a canopy over clover seedlings. Treatment may then be made up until grain is 8 to 10 inches tall (in tiller to early boot stage). With certain weed species that are more difficult to control and under conditions where normally easy to control

species are more resistant to weed killers, as much as 3/4 to 1 pint of AMOCO MCPA Weed Killer M may be required for effective weed control. Do not use on small grain interplanted with sweet clover or birdsfoot trefoil unless injury to the legume can be tolerated. Use on grain interplanted with seedling alfalfa only where specifically recommended by local Extension Service Weed Specialist or Agricultural Experiment Station. MCPA is safer on oats than 2,4-D.

### WEED CONTROL IN ESTABLISHED ALFALFA AND RED CLOVER

For control of Yellow Rocket and other susceptible winter annual weeds use 1 pint of AMOCO MCPA Weed Killer-M in the amount of water required for good coverage. Application must be made in very early spring before growth of the alfalfa or clover has commenced. Temperature at time of spraying should be above 40°F.

USDA REG. NO. 145-127  
(1971)

145-M DB 100-1  
MADE IN U.S.A.

ACCEPTED

FEB 8 - 1971

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 145-127

®

-M

INGREDIENTS BY WEIGHT  
Sodium salt of 2 Methyl  
chlorophenoxyacetic acid (MCPA  
(47.5%)) 52.1%

INGREDIENTS BY WEIGHT  
Water 47.9%

AMOCO MCPA Weed Killer M  
is equivalent of 4 pounds of  
2 Methylchlorophenoxyacetic acid

55 POUNDS