



Controls Horn Flies, Stable Flies,
Face Flies, House Flies, and Lice.

PRECAUTIONS IN USING

Harmful if swallowed. Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. Keep away from open flame. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water, or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Do not re-use container. Destroy when empty.

NON-WARRANTY NOTICE

Follow directions carefully. Seller warrants such directions to be proper and that product conforms to the descriptions on this label. Timing and method of application, weather and crop conditions, and other influencing factors in the use of the product are beyond the control of the Seller, so Seller makes no other warranty, express or implied (including a warranty of merchantability), with reference to the results or effects of use. No person has authority to make any representation not contained on this label.

EPA REG NO 1145-139
(043)


MADE IN U.S.A.

Bovinol

Ciovap*

A CIODRIN* AND VAPONA* INSECTICIDE
READY-TO-USE FLY SPRAY
FOR DAIRY AND BEEF CATTLE

WARNING: KEEP OUT OF THE REACH
OF CHILDREN. SEE OTHER WARNINGS
ON SIDE PANEL.



ACTIVE INGREDIENTS BY WEIGHT	
Dimethyl Phosphate of Alpha-Methylbenzyl	1.00%
3-Hydroxy-Cis-Crotonate**	0.23%**
2,2-Dichlorovinyl Dimethyl Phosphate	0.02%***
Related Compounds	98.75%
INERT INGREDIENTS BY WEIGHT	
TOTAL	100.00%

**CIODRIN Insecticide
***Equivalent to 0.25% VAPONA Insecticide

SEE USE AND DIRECTIONS FOR APPLICATION ON SIDE PANEL

GENERAL DIRECTIONS

BOVINOL CIOVAP Insecticide is a specially prepared ready-to-use insecticide suitable for use in conventional hand, power, or automatic spray equipment and in back rubbers. BOVINOL CIOVAP is approved for lactating dairy animals. No feed-off period is necessary when used on beef animals. Apply

with hand, power, or automatic equipment to moisten the hair only and not in excess of 2 fluid ounces daily per animal. Do not wet the skin due to possible skin irritation. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

For the control of certain insects attacking animals, apply in accordance with State and Federal Recommendations.

CONTENTS

GALLONS

PRODUCT OF AMOCO OIL COMPANY
CHICAGO, ILLINOIS, USA 60680

USE AND DIRECTIONS FOR APPLICATION

Lactating Dairy and Beef Animals: Flies (Horn Fly, Stable Fly, Face Fly, House Fly)—For the control of Horn Flies, Stable Flies, Face Flies, and House Flies, make a light application of BOVINOL CIOVAP daily as a fine mist to thoroughly cover all parts of the animal, including the legs.

Lactating Dairy and Beef Animals: Lice—Make 1 to 2 applications of 3 to 3½ ounces of BOVINOL CIOVAP at 14 day intervals as a fine mist spray to thoroughly cover all parts of the animal. If material is applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

Back and Face Rubbers: For the control of Horn Flies and Face Flies on lactating dairy and beef animals, use BOVINOL CIOVAP in back rubbers and face rubbers. Use 1 gallon for each 20 linear feet of burlap back rubber. For automatic back rubbers recharge the reservoir as needed. Animals must rub face on rubbers for best face fly control.

Do not treat animals within one day of slaughter.

Do not apply dermally in combination with Trichlorofon.

*SHELL REGISTERED TRADEMARK
774-OX-MP-L-0373



Bovinol

Ciovap*

A CIODRIN* AND VAPONA* INSECTICIDE
READY-TO-USE FLY SPRAY
FOR DAIRY AND BEEF CATTLE

WARNING: KEEP OUT OF THE REACH
OF CHILDREN. SEE OTHER WARNINGS
ON SIDE PANEL.

ACTIVE INGREDIENTS BY WEIGHT

Dimethyl Phosphate of Alpha-Methylbenzyl	
3-Hydroxy-Cis-Crotonate**	1.00%
2,2-Dichlorovinyl Dimethyl Phosphate	0.23%***
Related Compounds	0.02%***
INERT INGREDIENTS BY WEIGHT	98.75%
TOTAL	100.00%

**CIODRIN Insecticide

***Equivalent to 0.25% VAPONA Insecticide

SEE USE AND DIRECTIONS

with hand, power, or automatic equip-

USE AN
FOR AI

Lactati
Flies (F
Fly, Ho
Horn Flie
House F
BOVINC
to thorc
animal, i

Lactati
Lice — M
3½ ounce
day inte
thorough
If materi
ing equi
time per

Back a
control c
lactating
BOVINC
and face
20 linea
For aut
the rese
rub face