RYNOTOX®

ACTIVE INGREDIENT:

Ryanodine Alkaloid 0.08%
INERT INGREDIENTS 99.92%

USE: For use on Citrus Trees for the control of Citrus Thrips.

114818

SUGGESTED USES

Ground Rigs: Use RYNO-TOX at rate of 10-15 lbs. per acre in 20-100 gals, of water. Additional sugar may be added at the rate of 4-9 lbs. per acre. Do not overdilute. If 200 gallons of spray solution are applied per acre, increase dosage to 20 lbs. per acre and add extra sugar.

Air Application: Apply 10-20 lbs. per acre in 15-25 gallons of water with 3 ounces of TP!-SPRED™ A, or 1-2 pints per acre of SpreaderSticker 3-S. Additional sugar may be added at the rate of 4-9 lbs. per acre if desired.

This product contains 6 lbs. of Sugar for each 10 lbs.

NOTE: Use 20 mesh screen size or larger. Small screens may cause clogging.

CAUTION: Do not leave in water solution longer than necessary to spray out tank.

This product is toxic to fish.

Keep out of lakes, streams, or ponds.

Do not apply where runoff is likely to occur.

Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

CAUTION

KEEP OUT OF REACH OF CHILDREN HARMFUL IF SWALLOWED

AVOID INHALATION OF SPRAY MIST

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURY-ING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

EPA REG. NO. 1148-18-AA

(B) 1634

LEFFINGWELL CHEMICAL COMPANY BREA, CALIFORNIA